

Review

Key mediators modulating TAG synthesis and accumulation in woody oil plants

Yicun Chen¹, Yangdong Wang^{1*}, Guan Zhou^{1,2}, Peng Li^{1,2} and Shanshan Zhang^{1,2}

¹Institute of Subtropical Forestry, Chinese Academy of Forestry, Fuyang, 311400, China.

²College of Life Science, Anhui Normal University, Wuhu, 241000, China.

Accepted 27 November, 2008

Woody oil plant is gaining increasing interest as substitute for petroleum-derived materials, and its enriched hydroxy and conjugated fatty acids for industrial applications. In pursuit of high-value and level oils, a better understanding of mechanisms regarding triacylglycerol (TAG) synthesis and accumulation is required. Here we first summarized the germplasm resources of woody oil plants, and the key mediators on TAG synthesis and accumulation, among which diacylglycerol acyltransferases (DGATs) is discussed for its clear role in TAG amount and composition. Furthermore TAG-associated proteins called oleosins are also discussed in depth due to their determination on the amount and size of oil bodies. Previously, two transcripts of oleosins were isolated from *Vernicia fordii* by us, and the homology of oleosins is analyzed. TAG accumulation is the result of multi-play action of the above mediators at varied levels. Furthermore, the channel of fatty acids flux also serves as a limiting factor for determination of the rate of TAG accumulation. The knowledge of key mediators modulating TAG accumulation will provide new insight on further metabolic engineering of oilseeds in woody plants.

Key words: Woody oil plant, triacylglycerol accumulation, diacylglycerol acyltransferases, oleosin, fatty acid flux.

INTRODUCTION

As a result of reducing petrochemical resources and environmental consciousness, there will be a worldwide increasing demand for oil plants and their derivatives as substitutes for petrochemicals in industrial applications, such as biolubricants, biofuels, nylon precursors and detergent feedstocks (Metzger and Bornsheuer, 2006). According to international biodiesel standard, oils from woody oil plants have been defined as biodiesel materials, such as oils from *Camellia oleifera*, *Jatropha curcas*, *Elaeis guineensis*, etc. and others supply natural oil for industrial usage, such as *V. fordii*, *Keteleeria evelyniana*, *Pinus koraiensis*, etc. So far, tremendous application on crops and model plants has been performed for the production of biofuel feedstocks and oleo-chemicals. For example, soybean seeds are currently used for biodiesel production (Hill et al., 2006). Compared with oil crops, the woody oil plants relatively hold considerable superiority in two aspects. Firstly, wood

oil plants will not have competition with foodstuff and have no danger about food contamination. Secondly, woody plants enriched oils can produce novel fatty acid for industrial usage and feedstocks for biodiesel. However, the woody plants generally have limiting agronomic traits such as low yields, toxicity and the very limited geographical growing areas. Fortunately, genetic engineering offers an exciting opportunity. With a particular focus on woody oil plants oilseed, we review the woody oil plant resources and the key mediators on TAG complex process of synthesis and accumulation. It is well-known that three biosynthetic events are involved in the production of seed storage oils. The first is the fatty acids synthesis in plastids. The second refers to these fatty acids modification by enzymes located in the endoplasmic reticulum (ER). The third is the packaging of the fatty acids into TAG, which subsequently accumulate in oil bodies that bud off from the ER. The paper sets out to discuss the complex process of TAG synthesis and accumulation in woody plants at molecular level. In addition, environmental effect and carbon supply on accumulation were also taken into account.

*Corresponding author. E-mail: wyd11111@126.com.

WOODY OIL PLANTS ARE ATTRACTIVE FOR ITS USAGE

Woody oils plants are utilized for many food and industrial applications. They include edible oils, processed ingredients for the food industry and feedstocks for chemicals such as formulation of paints, inks, resins, vanish, plastics and biodiesel production. Table 1 summarized woody oil plants resources in the world.

KEY ENZYMES ON NOVEL FATTY ACID MANIPULATION IN WOODY OIL PLANTS

The constituent fatty acids in TAG confer specificity and selectivity of oil usage. Through metabolic engineering of the fatty acid composition, it is possible for woody oil plants to supply petroleum-derived materials in fuels, lubricants, and special chemicals. After fatty acids synthesis, there is an important process of manipulation, in which desaturases are active. They generally include Δ^9 and Δ^{12} desaturase and their various homologs. Recent researches have been focused on divergent forms of the Δ^{12} -oleic acid desaturase (FAD2-like enzyme). Divergent FAD2 forms catalyzed a wide range of fatty acid modification, including conjugation, hydroxylation and epoxygenation (Iwabuchi et al., 2003; Cahoon and Kinney, 2005). For instance, α -eleostearic acid (18:3 $\Delta^{9cis, 12cis, 15cis}$) in *V. fordii* (tung tree) is catalyzed by Δ^{12} fatty acid conjugase FADX (Dyer et al., 2002). FAD2-like conjugase from *punica granatum* (pomegranate) for punicic acid has also been described (Hornung et al., 2002; Iwabuchi et al., 2003). Furthermore, a desaturase/hydroxylase bifunctional enzyme from *Lesquerella fendleri* has been identified to produce lesquerolic acid (Broun et al., 1998; Moon et al., 2001).

MEDIATORS ON TAG SYNTHESIS AND ACCUMULATION IN WOODY OIL PLANTS

The main lipids store in plants are TAGs. Generalized pathways of TAG synthesis and key mediators on its accumulation especially in the woody oil plants are depicted in this section. Our current investigation on tung tree is also mentioned here.

Pathways leading to TAG formation

Formation of TAG can be achieved in several ways, as shown in Figure 1. The first classical Kennedy pathway (Kennedy, 1961), using acyl-CoA as donor and diacylglycerol (DAG) as acceptor, is catalyzed by enzyme acyl-CoA: diacylglycerol acyltransferase (DGAT) and transfers an acyl group from acyl-CoA to sn-3 of DAG and forms TAG. The second is acyl-CoA independent pathway, which uses phosphatidylcholine (PC) as acyl

donors and DAG as acceptor, catalyzed by an enzyme called phospholipids: diacylglycerol acyltransferase (PDAT), which can transfer the sn-2 acyl chain from PC to DAG, forming lyso-PC and TAG (Dahlqvist et al. 2000). Also the acyl-CoA independent pathway can be catalyzed by DGAT, using two molecules of DAG to produce TAG and monoacylglycerol (MAG) (Weselake, 2005).

Key mediators involved in TAG biosynthesis and manipulation

Much more attention has been focused on the enzymes involved in the TAG synthesis: phospholipase A₂ (PLA₂) and G-3-P acyltransferase (GPAT), as well as phospholipids: diacylglycerol acyltransferase (PDAT), Lyso PA acyltransferase (LPAAT) and diacylglycerol acyltransferase (DGAT). Currently, at least eight GPAT genes have been identified in *Arabidopsis thaliana*, but neither of them has been shown important in seed TAG biosynthesis (Beisson et al., 2007). PDAT catalyses the acyl transfer from PC to SN-1,2-DAG to yield TAG. But little evidence has been found to date that it plays a major quantitative or qualitative role in seed TAG metabolism (Dahlqvist et al., 2000). LPAAT genes have been proven to be useful for increasing the accumulation of target fatty acids in TAG in transgenic crops. Scarce information of these genes in woody plant has been obtained. Considering the development of woody oil plants seeds metabolic engineering, knowledge-based gene discovery and application are urgently required.


Now there is general agreement that diacylglycerol acyltransferases (DGATs) exert a strong influence on the amount and composition of TAG synthesized in developing seeds. DGAT catalyzed the committed step of oil biosynthesis by transferring a fatty acyl group from acyl-CoA to a diacylglycerol substrate to form TAG (Figure 1), and can also could employ two molecules of DAG to produce TAG and monoacylglycerol (MAG). At least three different and structurally unrelated enzymes catalyze DGAT activity, including DGAT1, DGAT2 (Shockey et al., 2006) and DGAT3 (Saha et al. 2006). For woody oil plants, DGAT1 and DGAT2 have been investigated in Tung tree, where DGAT2 is strongly induced in developing seeds, and the timing of DGAT2 gene expression coincides closely with the onset of eleostearic acid biosynthesis and total oil accumulation. Expression of tung tree DGAT2 in yeast cells resulted in elevated accumulation of TAG than DGAT1 (Shockey et al., 2006). Currently, it has been reported that DGAT1-2 with F469 a phenylalanine insertion is responsible for the increased oil and fatty acid contents in maize (Zheng et al., 2008). Lardizabal et al. (2008) showed that expression of a codon-optimized version of DGAT2A from soil fungal *Umbelopsis ramanniana* in soybean resulted in oil increase in seeds (Lardizabal et al., 2008), which offered an original indication for DGAT transgenic engineering. In addition to catalyzing a critical role in TAG synthesis,

Table 1. Woody oil plants resources summarization and their oil amounts.

Woody plant species	Oil position	Oil (%)	Woody plant species	Oil position	Oil (%)
<i>Aconitum flavum</i>	seed	46.9	<i>Lasiococca comberi</i>	seed	59.3
<i>Actinodaphne obovata</i>	kernel	51	<i>Lindera aggregata</i>	seed	53.1
<i>Aleurites moluccana</i>	kernel	65.4	<i>Lindera caudata</i>	Fruit	50.5
<i>Amesiodendron chinense</i>	kernel	50.9	<i>Lindera chienii</i>	seed	49.3
<i>Anacardium occidentale</i>	kernel	50.1	<i>Lindera chunii</i>	seed	54.1
<i>Arachis hypogaea</i>	kernel	50.7	<i>Lindera communis</i>	seed	53.1
<i>Butyrospermum parkii</i>	kernel	52.5	<i>Lindera latifolia</i>	seed	57.6
<i>Carya cathayensis</i>	kernel	67.1	<i>Lindera megaphylla</i>	seed	52.2
<i>Carya hunanensis</i>	kernel	54.8	<i>Lindera metcalfiana</i>	seed	57.3
<i>Carya illinoensis</i>	kernel	63.5	<i>Lindera nacusua</i>	kernel	61.6
<i>Camellia furfuracea</i>	kernel	52.1	<i>Lindera reflexa</i>	seed	52.5
<i>Camellia microphylla</i>	kernel	59.2	<i>Lindera thomsonii</i>	seed	50.5
<i>Camellia obtusifolia</i>	kernel	50.5	<i>Litsea chunii</i>	seed	49.4
<i>Camellia oleifera</i>	kernel	49.4	<i>Litsea coreana</i>	seed	61.9
<i>Camellia pitardii</i>	kernel	56	<i>Litsea cubeba</i>	seed	49.1
<i>Camellia reticulata</i>	kernel	58.3	<i>Litsea elongata</i>	seed	53
<i>Camellia sasanqua</i>	kernel	47.6	<i>Litsea euosma</i>	kernel	56.2
<i>Camellia semiserrata</i>	kernel	63	<i>Litsea glutinosa</i>	kernel	57.5
<i>Camellia vietnamensis</i>	kernel	48.1	<i>Litsea panamonja</i>	kernel	51.2
<i>Canarium album</i>	kernel	58.1	<i>Litsea populifolia</i>	Fruit	49.4
<i>Canarium bengalense</i>	kernel	57	<i>Litsea pungens</i>	kernel	55.4
<i>Canarium pimela</i>	kernel	59.4	<i>Litsea rotundifolia</i>	seed	62.5
<i>Celtis wightii</i>	kernel	68.1	<i>Macaranga adenantha</i>	kernel	60.3
<i>Cephalotaxus fortunei</i>	kernel	63.6	<i>Macadamia ternifolia</i>	kernel	66.5
<i>Cephalotaxus sinensis</i>	kernel	63.3	<i>Madhuca pasquieri</i>	kernel	46.6
<i>Cerbera manghas</i>	kernel	59.1	<i>Malania oleifera</i>	kernel	58.2
<i>Cinnamomum burmannii</i>	seed	59.4	<i>Maytenus austroyunnanensis</i>	seed	57.4
<i>Cinnamomum glanduliferum</i>	kernel	59.9	<i>Maytenus hookeri</i>	seed	56.6
<i>Cinnamomum japonicum</i>	seed	58.3	<i>Mesua nagassarium</i>	kernel	76.5
<i>Cinnamomum porrectum</i>	kernel	55.4	<i>Melliodendron xylocarpum</i>	kernel	49.6
<i>Cinnamomum saxatile</i>	kernel	54.8	<i>Momordica macrophylla</i>	kernel	46.4
<i>Cinnamomum septentrionale</i>	kernel	56.3	<i>Neocinnamomum caudatum</i>	kernel	57.4
<i>Citrus grandis</i>	kernel	49.5	<i>Neocinnamomum delavayi</i>	kernel	62.8
<i>Citrus reticulata</i>	kernel	46.8	<i>Neolitsea aurata</i>	seed	54.1
<i>Cocos nucifera</i>	fruit	59.3	<i>Neolitsea phanerophlebia</i>	Fruit	51.4
<i>Cordia dichotoma</i>	seed	51.8	<i>Neolitsea umbrosa</i>	Fruit	57.5
<i>Corylopsis multiflora</i>	kernel	51.5	<i>Ostodes paniculatus</i>	kernel	55
<i>Corylus ferox</i>	fruit	62.9	<i>Persea americana</i>	fruit	50
<i>Corylus heterophylla</i>	fruit	54.4	<i>Pinus armandi</i>	kernel	57.1
<i>Corylus mandshurica</i>	fruit	63.8	<i>Pinus koraiensis</i>	kernel	69.7
<i>Delavaya yunnanensis</i>	kernel	70	<i>Pistacia chinensis</i>	Fruit	24.4
<i>Deutzianthus tonkinensis</i>	kernel	49.7	<i>Polyalthia plagioneura</i>	kernel	54.7
<i>Diplopanax stachyanthus</i>	kernel	57.1	<i>Podocarpus nagi</i>	kernel	48.4
<i>Dracontomelon duperreanum</i>	kernel	64	<i>Prunus amygdalus</i>	kernel	58.4
<i>Dracontomelon macrocarpum</i>	kernel	69.5	<i>Prunus armeniaca</i>	kernel	51.9
<i>Dysoxylum binectariferum</i>	Fruit	52.2	<i>Prunus davidiana</i>	kernel	50.9
<i>Eberhardtia aurata</i>	kernel	51.3	<i>Prunus mira</i>	kernel	50.6
<i>Eberhardtia tonkinensis</i>	kernel	57.5	<i>Prunus sibirica</i>	kernel	49.9
<i>Elaeis guineensis</i>	fruit	65.6	<i>Prunus undulata</i>	kernel	54.2
<i>Euonymus acanthocarpus</i>	seed	47	<i>Pterospermum menglungense</i>	seed	47.6

Table 1. Contd.

<i>Euonymus carnosus</i>	seed	47.8	<i>Pyralaria edulis</i>	kernel	62.5
<i>Euonymus grandiflorus</i>	seed	51.2	<i>Pyralaria sinensis</i>	kernel	60.7
<i>Euonymus hamiltonianus</i>	seed	52.6	<i>Sarcandra glabra</i>	seed	51.2
<i>Euonymus laxiflorus</i>	seed	46.3	<i>Santalum album</i>	kernel	62.6
<i>Euonymus macropterus</i>	seed	47.5	<i>Sapium japonicum</i>	seed	54.7
<i>Euonymus myrianthus</i>	seed	49.3	<i>Schleichera trijuga</i>	kernel	68.8
<i>Euonymus streptopterus</i>	seed	47.4	<i>Scleropyrum wallichianum</i>	kernel	66.4
<i>Fagus longipetiolata</i>	seed	42.8	<i>Sesamum indicum</i>	seed	51.1
<i>Handeliodendron bodinieri</i>	kernel	52.6	<i>Sloanea hemsleyana</i>	seed	51
<i>Hodgsonia macrocarpa</i>	kernel	64.2	<i>Sloanea sinensis</i>	seed	49.5
<i>Horsfieldia pandurifolia</i>	kernel	56.2	<i>Styrax confusus</i>	kernel	52.8
<i>Hydnocarpus</i>	kernel	51.2	<i>Sumbaviopsis albicans</i>	kernel	58.2
<i>Juglans cathayensis</i>	kernel	68.6	<i>Thevetia peruviana</i>	kernel	72.4
<i>Juglans mandshurica</i>	kernel	68.2	<i>Treminalia catappa</i>	kernel	59.4
<i>Juglans regia</i>	kernel	66.2	<i>Trewia nudiflora</i>	kernel	57.9
<i>Juglans sigillata</i>	kernel	69.4	<i>Tutcheria championi</i>	kernel	59.1
<i>Jatropha curcas</i>	kernel	56.6	<i>Tutcheria hirta</i>	kernel	64.9
<i>Keteleeria davidiana</i>	seed	47.9	<i>Vernicia montana</i>	kernel	52.5
<i>Keteleeria evelyniana</i>	seed	59.7	<i>Vernicia fordii</i>	kernel	80.0
<i>Keteleeria fortunei</i>	seed	48.4	<i>Xanthoceras sorbifolia</i>	kernel	59.9






Figure 2. The homology of oleosins in woody plants. There is conserved "proline knot": PX₅SPX₃P-. The oleosin of *V. fordii* is mostly homologous to *Jatropha curcas* and *C. oleifera*.

DGAT also has been shown involved in lipid metabolism associated with germination and leaf senescence. Furthermore, DAGT2 has been reported as a key regulator on solving the "bottleneck" (Figure 1), that is, the low TAG production in transgenic plants (Cahoon et al., 2007). In a word, the work above highlights DGATs as promising targets for manipulating TAG content and the constituent fatty acids.

Oleosins - the proteins could modulate oil body structure and TAG accumulation

TAGs accumulate in structures known as oil bodies or oleosomes, the number and size of which are correlative with TAG content. Oil bodies vary in diameter from about 0.6 to 2 μm (Huang, 1992) and comprise a matrix of TAG surrounding by a phospholipids monolayer embedded by one major type of proteins called oleosins, which are usually presented as two or more isoforms and mainly expressed in seeds. Oleosins share similar structure

properties including a long hydrophobic domain around a 12-amino acid motif called a proline knot (Abell et al., 1997), both of which are essential for correct targeting to oil bodies (Van Rooijen and Moloney, 1995). Oleosins promote steric hindrance and electrical repulsion between oil bodies and prevent the organelles from coalescing during seed maturation, desiccation and germination (Tzen et al., 1992; Leprince et al., 1998). Oleosins accumulation has been proved to determine the size of oil bodies through the experiment that oleosin suppression resulted in an aberrant phenotype of embryo cells that contain unusually large oil bodies (Siloto et al., 2006). We first isolated oleosin cDNAs from *V. fordii*; they are most homologous to those in *J. curcas* and *C. oleifera* in oil woody plants, as indicated in Figure 2.

Moreover, the suppression of oleosins had a small but statistically significant effect on fatty acid preferences in TAG (Siloto et al., 2006). Therefore, the introduction of a foreign oleosin may be an alternative way to choose fatty acid and increase TAG accumulation through modulating oil body size.

Direction of fatty acids flux into TAG

The production of high TAG level enriched novel fatty acid is the pursuit of biologists. But most of the cases are undesirable especially in transgenic plants. The exotic fatty acids generally could not be stored in the right place. Most of them are integrated in phosphatidylcholine (PC) instead of storing in oil bodies and ultimately in TAG form, depicted as Figure 1B. For example, FAD2-like genes isolated from *Morordica charantia* and *Impatiens balsamina* produce α -eleostearic acid and α -parinaric acid in transgenic plants only at a third of the level in the native species (Cahoon et al., 1999). The results strongly indicated that there is a fatty acid channel into TAG after synthesis. The direction of fatty acids flux between the PC, acyl-CoA and TAG pools determines final TAG level (Lardizabal et al., 2008) (Figure 1A). The above key mediators in TAG synthesis, including GPAT, LPAAT, DGAT and PDAT, seemed to be involved in prevention of the unusual fatty acids in PC and other membrane phospholipids and directs them into TAG. Among them, DGATs enzymes are proven to be more crucial for the direction of fatty acid flux into TAG (Zheng et al., 2008; Shockey et al., 2006; Kroon et al., 2006).

CONCLUSION AND PERSPECTIVES

Woody oil plants, as a substitute of oil crops to supply petroleum-derived materials and wide application in industry, have a prominent future. However, large-scale production of these woody oil plant species through traditional farming and breeding procedure has a diversity of problems, such as poor agronomic traits and management. With the fast development of molecular biology and the study on the model plants, the most promising route to the development of such renewable resources is via oilseed genetic engineering. Recently, knowledge about TAG synthesis and accumulation in woody oil plants has been gradually realized. Tung tree for example, FAD2, FAX, Oleosins, DGAT1 and DGAT2 has been well studied, but for the woody oil plants, there is still a long way to go to obtain useful information.

In view of complicated processes participating in TAG synthesis and accumulation, the three aspects should be taken into account in increasing TAG accumulation. The first is that the multi-play and multi-level regulation should be elucidated. In transgenic plants, the mediators always cooperate at transcriptional and translational level. The second is the cellular regulation of mediators. The compartmentalization of enzymes activity within specific regions or subdomains of the ER might be essential for both novel fatty acid synthesis and their channeling into TAG (Dyer and Mullen, 2008). The biochemical and immunolocalization studies of FAD2-like enzyme, such as fad2 and FADX in tung tree, have shown their location exclusively in the ER and this introduces a sort of topo-

logical orientation in ER membranes (Dyer et al. 2002). DGATs are also localized in subdomains of ER and hold potential for compartmentalization of oil biosynthesis (Shockey et al., 2006; Dyer and Mullen, 2008). The third aspect is that the environmental factors including light intensity, temperature, mineral deprivation, osmotic agent, together with alteration of carbon source also contribute to the process of TAG accumulation and so should be considered in farming and management.

REFERENCES

- Abell BM, Holbrook LA, Abenes M, Murphy DJ, Hills MJ and Moloney MM (1997). Role of the proline knot motif in oleosin endoplasmic reticulum topology and oil body targeting. *Plant Cell*. 9: 1481-149.
- Beisson F, Li Y, Bonaventure G, Pollard M, Ohlrogge JB (2007). The acyltransferase GPAT required for the synthesis of suberin in seed coat and root of *Arabidopsis*. *Plant Cell*. 19: 351-368.
- Broun P, Boddupalli S, Somerville C (1998). A bifunctional oleate 12-hydroxylase: desaturase from *Lesquerella fendleri*. *Plant J*. 13: 201-210.
- Cahoon EB, Carlson TJ, Ripp KG, Schweiger BJ, Cook GA, Hall SE, Kinney AJ (1999). Biosynthetic origin of conjugated double bonds: Production of fatty acid components of high-value drying oils in transgenic soybean embryos. *Proc. Natl. Acad. Sci*. 96: 12935-12940.
- Cahoon EB, Kinney AJ (2005). The production of vegetable oils with novel properties: using genomic tools to probe and manipulate plant fatty acid metabolism. *Eur. J. Lipid. Sci*. 107: 239-243.
- Cahoon EB, Shockey JM, Dietrich CR, Gidda SK, Mullen RT, Dyer JM (2007). Engineering oilseeds for sustainable production of industrial and nutritional feedstocks: solving bottlenecks in fatty acid flux. *Curr. Opin. Plant Biol*. 10: 236-244.
- Dahlqvist A, Ståhl Ulf, Lenman M, Banas A, Lee M, Sandager L, Ronne H, Szymne S (2000). Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes the acyl-CoA independent formation of triacylglycerol in yeast and plants. *Proc. Natl. Acad. Sci*. 97: 6487-6492.
- Dyer JM, Chapital DC, Kuan JCW, Mullen RT, Turner C, Mckeon TA, Pepperman AB (2002). Molecular analysis of a bifunctional fatty acid conjugase/desaturase from Tung. implications for the evolution of plant fatty acid diversity. *Plant Physiol*. 130: 2027-2038.
- Dyer JM, Mullen RT (2008). Engineering plant oils as high-value industrial feedstocks for biorefining: the need for underpinning cell biology research. *Physiol. Plant*. 132: 11-22
- Hill J, Nelson E, Tilman D, Polasky S, Tiffany D (2006). Environmental, economic and energetic costs and benefits of biodiesel and ethanol biofuels. *Proc. Natl. Acad. Sci*. 103: 11206-11210.
- Hornung E, Pernstich C, Feussner I (2002). Formation of conjugated Delta11Delta13-double bonds by Delta12-linoleic acid (1,4)-acyl-lipid-desaturase in pomegranate seeds. *Eur. J. Biochem*. 269: 4852-4859.
- Huang AHC (1992). Oil bodies and oleosins in seeds. *Ann. Rev. Plant Physiol. Plant Mol. Biol*. 43: 177-200.
- Iwabuchi M, Kohno-Murase J, Imamura J (2003). 12-Oleate Desaturase-related enzymes associated with formation of conjugated *trans*-11, *cis*-13 Double Bonds. *J. Biol. Chem*. 278: 4603-4610.
- Kennedy EP (1961). Biosynthesis of Complex Lipids, *Fed. Proc. Am. Soc. Exp. Biol*. 20: 934-940.
- Kroon JTM, Wei W, Simon WJ, Slabas AR (2006). Identification and functional expression of a type 2 acyl-CoA:diacylglycerol acyltransferase (DGAT2) in developing castor bean seeds which has high homology to the major triglyceride biosynthetic enzyme of fungi and animals. *Phytochemistry*. 67: 2541-2549
- Lardizabal K, Effertz R, Levering C, Mai J, Pedroso MC, Jury T, Aasen E, Gruys K, Bennett K (2008). Expression of *Umbelopsis ramanniana* DGAT2A in seed increases oil in soybean. *Plant Physiology*. in press
- Leprince O, Van Aelst AC, Pritchard HW, Murphy DJ (1997). Oleosins prevent oil-body coalescence during seed imbibition as suggested by

- a low-temperature scanning electron microscope study of desiccation-tolerant and -sensitive oilseeds. *Planta*, 204: 109-119
- Metzger JO, Bornsheuer U (2006). Lipids as renewable resources: current state of chemical and biotechnological conversion and diversification. *Appl. Microbiol. Biotechnol.* 71: 13-22.
- Moon H, Smith MA, Ljerka K (2001). A condensing enzyme from the seeds of *Lesquerella fendleri* that specifically elongates hydroxy fatty acids. *Plant Physiol.* 127: 1635-1643.
- Saha S, Enugutti B, Rajakumari S, Rajasekharan R (2006). Cytosolic triacylglycerol biosynthetic pathway in oilseeds. Molecular cloning and expression of peanut cytosolic diacylglycerol acyltransferase. *Plant Physiol.* 141: 1533-1543.
- Shockey JM, Gidda SK, Chapital DC, Kuan JC, Dhano PK, Bland JM, Rothstein SJ, Mullen RT, Dyer JM (2006). Tung tree DGAT1 and DGAT2 have nonredundant functions in triacylglycerol biosynthesis and are localized to different subdomains of the endoplasmic reticulum. *Plant Cell.* 18: 2294-2313.
- Siloto RMP, Findlay K, Lopez-Villalobos A (2006), Yeung, EC, Nykiforuk CL and Moloney MM. The accumulation of oleosins determines the size of seed oilbodies in *Arabidopsis*. *Plant Cell.* 18: 1961-1974.
- Tzen JTC, Lie GC, Huang AHC (1992). Characterization of the charged components and their topology on the surface of plant seed oil bodies. *J. Biol Chem.* 267: 15626-15634.
- Van Rooijen GJH, Moloney MM (1995). Structural requirements of oleosin domains for subcellular targeting to the oilbody. *Plant Physiol.* 109: 1353-1361.
- Weselake RJ (2005). Storage lipids in: Murphy, Denis J. ed. *Plant Lipids - biology, utilization and manipulation*. Blackwell, Oxford, p: 162-225
- Zheng P, Allen WB, Roesler K, Williams ME, Zhang S, Li J, Glassman Kimberly, Ranch J, Nubel D, Solawetz W, Bhatramakki D, Llaca V, Deschamps S, Zhong GY, Tarczynski MC, Shen B (2008). A phenylalanine in DGAT is a key determinant of oil content and composition in maize. *Nat. Genet.* 40: 367-372.