

Full Length Research Paper

Lead and zinc concentrations in hair and nail of some Kano inhabitants

J. T. Ayodele* and A. S. Bayero

Department of Chemistry, Bayero University, P.M.B 3011, Kano-Nigeria.

Accepted 8 June, 2009

The advantages of hair and nail tissues analysis over other biological samples are that trace metal concentrations in them are not subjected to rapid fluctuation due to diet, air and water hence a long term stable nutritional status. Lead and zinc concentrations in hair and nail samples were determined by flame atomic absorption spectrophotometer (AAS). The mean zinc concentration in hair and nail were 0.695 ± 0.33 mg/g and 0.640 ± 0.52 mg/g respectively while the mean lead concentrations in hair and nail were 0.384 ± 0.34 and 0.463 ± 0.364 mg/g respectively. A progressive increase in zinc concentrations in hair and nails with age indicated no significant difference when their means were compared suggesting that zinc in hair and nails originate from a common source, comparing the mean lead concentrations in hair with the nails a significant difference is indicated in the 2 tissues ($p \leq 0.05$). Human hair and nails are therefore recording filaments that can reflect metabolic changes of many elements over long periods of time and hence furnish a print out of post nutritional event as dietary levels of some of the essential micro-elements.

Key words: Lead, zinc, hair, nail, AAS, Kano, Nigeria.

INTRODUCTION

The results of analysis of human milk, urine, saliva and sweat reflect some of the components that are absorbed but excreted from the body. The blood contains components absorbed and temporarily in circulation before excretion and/or storage (EPA, 1980). The hair, nails and teeth tissues in which trace minerals are sequestered and/or stored and can be used to effectively monitor the highest priority toxic trace metals (Barrett, 1985; Afridi et al., 2006a, b; Kazi et al., 2008). Hair and nails are recording filaments that can reflect metabolic changes of many elements over long periods of time which may furnish post nutritional events (Strain et al., 1966, 1972; Hambidge, 1982; Klevay, 1997).

Analysis of trace and heavy metal in hair and nail is a simple laboratory test which helps to monitor how well bodies are responding to our diets and environment (EPA, 1979). The advantages of hair and nail tissues analysis over other diagnostic samples are, trace metal concentrations are not subjected to rapid fluctuation due to diet, air and water hence a long term stable nutritional status. Sample collection is non-evasive, stable at room temperature, analytical methods of analysis are easy as their con-

centrations in hair and nail are high compared to other measurements (Borel and Anderson, 1984; Bord and Anderson, 1984). With progress in measuring and understanding the specific functions of macro, trace and toxic elements in human physiology, it has become evident that the action of each element may be potentiated or reduced by the presence of another. This may be why the ratio between the concentrations of any given element in body chemistry is decisive as to whether or not deficiencies or toxicities may occur. Therefore the requirement and the nutritional adequacy of a particular element depend on others already present in the body chemistry (Hill and Matrone, 1970). Zinc is important in human physiology and its deficiency leads to poor growth, hypogonadism and reduced immunity. In children it is associated with autism, dyslexia, apathy, lethargy irritability and childhood hyperactivity. In adults it has been linked with the development of both senility and Alzheimer's disease (Tuormaa, 1995).

Zinc an essential element in animal nutrition is required in amounts less than 100 mg/kg in the dry matter (Hambidge et al., 1987; Underwood, 1977; Hill et al., 1987; Neathery et al., 1973). Zinc is an essential component of carbonic anhydrate (Todd et al., 1934; Hove et al., 1940). It is an indispensable component of over 200 enzymes or proteins (Hambidge et al., 1987). When in short supply the

*Corresponding author. E-mail: tjayodele@yahoo.com.




Figure 1. Frequency distribution pattern for AGE (years) of hair and nail donors.

following activities may be decreased, viz plasma alkaline phosphates, liver, retina and testes alcohol, dehydrogenate and connective tissue, thymidine kinase, pancreatic carboxyl peptidase A and liver nuclear DNA-dependent RNA-polymerase (Hambidge et al., 1987; Miller et al., 1979).

Zinc is involved in nucleic acid and protein metabolism and in the fundamental processes of cell differentiation and replication. It is chemically bound in crystalline insulin. It plays a role in the production, storage and release of several other hormones as well as in the effectiveness of receptor sites and end organs responsiveness (Hambidge et al., 1987). It is essential for maintaining normal growth, reproduction and lactation performance (Miller et al., 1979). It is associated with taste and smell acuity and wound and burn healing. It is essential to the integrity of the immune system as it plays a role in stabilization of cell membranes and microtubule polymerization (Hambidge et al., 1987).

Lead an ubiquitous element in biological samples does not possess any evidence suggesting its importance in normal metabolic processes in man. It has been implicated in red cell survival and in the synthesis of globin (Boeckx, 1986). Its level in human hair and nails are yet to be determined accurately (Tuormaa, 1994).

Realizing that toxic metal excesses and trace mineral deficiencies are associated with all forms of reproductive failures, the hair tissue analyses before conception has been advocated for (Barnes and Bradley, 1994; Bradley and Bennett, 1995). This paper reports the determination of lead and zinc in human hair and nails.

MATERIALS AND METHODS

Lead and zinc were determined from various volunteered subjects

resident in Kano over a 6 month period. 350 hair and 300 nail samples were collected from subjects in the age range of 1 - 55 years. Nail samples were collected in polyethylene containers and were washed in 1% solution of TRITONX-100 in de-ionized water in an ultra sonic bath and on drying were stored in small plastic tubes (Iyengar, 1984). Hair samples were collected from each subject as close to the scalp as possible (Kucera et al., 1996). To eliminate grease and surface contamination the hair samples were rinsed with acetone (Kucera et al., 1996) and by washing each sample in detergent and distilled water (Nowak, 1998; Martin et al., 2005) after which they were kept in an alcohol ether mixture for 45 mins and dried at 60°C for 72 h. 1.0 g of each sample was digested in 10 cm³ concentrated HNO₃ and the resulting solution was evaporated to dryness and redissolved in 0.1 M nitric acid. Trace metal concentrations were determined by flame atomic absorption on a model 210 VGP spectrophotometer attached to IBM personal computer. The result of the absorbance of each sample was the average of 10 sequential readings. Background light absorption and scattering were compensated for either by deuterium hollow cathode lamp or by tungsten/halogen lamp. Distilled water was digested as blank using the same procedure previously described (Ayodele and Abubakar, 1998; Ayodele and Abubakar, 2001)

Statistical analysis

All statistical computations either were on the PC 486 66 MHZ microcomputer using the integrated statistical package for windows from Umstat Ltd. (London) or dedicated micro instructions for the excel spread sheets from microsoft. The approach enabled the advantages of the various computational and graphical facilities of both types of software's to be used with the ability to read different file formats. The analyses of variance (ANOVA) were carried out according to the procedures described by O'Mahony (1986).

RESULTS AND DISCUSSION

The frequency distribution pattern for the age of hair and nail donors is as shown in Figure 1. The distribution is multimodal with a mean age of 27.51 ± 16.5 years. The frequency distribution pattern for zinc in hair and nail is as shown in Figure 2. This is multimodal for zinc in hair with a mean and standard deviation of 0.695 ± 0.33 mg/g whilst the frequency distribution pattern for zinc in nails of the sampled is bi modal and is skewed towards low frequency of higher concentrations with a mean and standard deviation of 0.64 ± 0.52 mg/g. The observed trends for zinc in hair and nails with respect to age of donors are as shown in Figure 3. Table 1 shows the maximum, minimum, mean, standard deviation and number of samples for zinc in hair and nails. Figure 3 summarizes zinc concentrations in hair and nails with respect to age of their donors. A progressive increase in zinc concentrations in hair and nails with age indicated no significance difference indicated when the mean zinc in hair was compared with that of the nail ($p < 0.05$). Our results are in agreement with several other authors who reported varying concentrations of these metals in hair and nail samples (Tables 2 and 3).

From the zinc levels in hair and nails it is reasonable to believe that zinc in human tissues may be playing some physiological roles (Vivoli et al., 1990). It is also reasonable to suggest that zinc in hair and nails originate from a number of sources, such as the air, water and food we con-




Figure 2. Frequency distribution pattern for zinc (mg/g) in hair and nails.




Figure 3. Zinc concentration in hair and nails with respect to age.

sume (Strain et al., 1972; Maugh, 1978; Casey and Hambidge, 1980). These high concentrations of zinc

secreted than could possibly be received from water, confirm the suspicion that there may be other sources since

Table 1. Lead and Zinc concentrations (mg/g) in hair and nails.

Concen.	Lead		Zinc	
	Hair	Nails	Hair	Nails
Maximum	1.48	1.35	1.33	1.90
Minimum	0.05	0.05	0.12	0.05
Mean	0.384	0.463	0.695	0.640
Std.Dev	0.34	0.364	0.33	0.52

Table 2. Results of zinc concentrations in hair and nails from different countries

Country	Mean	Units	References
Austria	174	µg/g	Ryabukin (1978)
Germany	129 (Hair) 108 (Nails)	µg/g µg/g	Wilhelm et al (1991)
New Jersey (USA)	205	µg/g	DeAntonio et al (1982)
Brazil	151- 168	µmol/g	Sandra et al (2002)
Saudi Arabia	121-247	µg/g	Imran et al (2003)
India	140.6±6.0	Ppm	Sukumar and Subramanian (2003)
India	110.29-286.59(nail)	µg/g	Mehra and Juneja (2005)
Turkey	21.40 – 176.96	Ppm	Ulvi et al (2003)
Poland	156.48± 74.5	mg/kg	Chojnacka et al(2005)
Poland	173-189	mg/kg	Chojnacka et al(2006)
Nigeria	57.7-510 (Hair)	µg/g	Nnorom et al(2005)
Nigeria	0.695±0.33 (Hair) 0.640 ±0.52 (Nail)	mg/g mg/g	This study

Table 3. Results of lead concentrations in hair from different countries.

Country	Mean	Range	Units	References
Austria		0.97-44.9	Ppm	Fergusson (1990)
Canada	10.1	0.5-25	ppm	Fergusson (1990)
Canada	16.90	0.5-35	ppm	Fergusson (1990)
Canada	6.3±0.90	10-350	ppm	Ferguson (1990)
India		10.40-67.00	µg/g	Sukumar and Subramanian (2003)
Indonesia	131.7±93.40		µg/g	Pirsaraei (2007)
Iran	21.1±13.20		ppm	Pirsaraei (2007)
New Zealand	14.50	14.42-48.30	µg/g µg/g	Fergusson (1990) Boris (1994)
Russia	6.55-16.20	6.41-7.37 6.55-16.2	µg/g	Boris (1994)
Saudi Arabia	1.046±1.39		mg.kg	Imran et al (2003)
U.S.A		7.6-107.1	ppm	Ferguson(1990)
Poland		1.42-2.17	ppm	Chojnacka et al (2005)
Poland		8.64-129.42	mg/kg	Chojnacka et al (2006)
India			µg/g	Mehra and Juneja (2005)
Nigeria	0.384±0.34 (Hair)	9.1-194.5	µg/g	Nnorom et al (2005)
Nigeria	0.464±0.364 (Nail)		mg/g mg/g	This study This study




Figure 4. Frequency distribution pattern for lead (mg/g) in hair and nails.




Figure 5. Lead in hair and nails with respect to age.

since the nutritional status of individual may contribute this effect (Calabrasc, 1980; Harding-Barlow, 1983; EPA, 1988; Ayodele and Bayero, 2002).

The frequency distribution pattern for lead in hair and nail tissues are as shown in Figure 4. The distributions are both skewed towards low frequencies of high concentrations with a mean and standard deviation of 0.384 ± 0.34 and 0.463 ± 0.364 mg/g in hair and nail respectively. Lead concentrations in hair and nail with respect to age of donors are as shown in Figure 5. Comparing hair and nails

as points of excretion the latter appear superior to the former. Comparing the mean lead concentrations in hair with the nails a significant difference is indicated in the lead concentrations in the 2 tissues ($p \leq 0.05$). It may thus be stated that human hair and nails are recording filaments that can reflect metabolic changes of many elements over long periods of time and hence furnish a print out of post nutritional event (Strain et al., 1972) as dietary levels of some of the essential micro-elements have been reported to correspond to hair concentrations of the elements (Rein-

hold et al., 1966; Strain et al., 1966; Potter et al., 1974; Maugh, 1978; Katz, 1979; Hopps, 1977; Casey and Hambidge, 1980).

Conclusion

Hair analysis provides a cosmetologist, a nutritionist and a doctor an additional diagnosis tool in their respective professional fields. The cosmetologists use this system to diagnose hair and skin related problems. If and when it is discovered that these problems are related to general health conditions they are referred to the appropriate professionals.

REFERENCES

- Afridi HI, Kazi TG, Jamali MK, Kazi GH, Arain MB, Jalbani N, and Memon AR (2006a). Atomic absorption spectrometric determination of Zn and Cr levels in scalp human hair samples. Influence of age gender and diabetic condition. *J. Indian Chem. Soc.* 83(10):1062-1064.
- Afridi HI, Kazi TG, Jamali MK, Kazi GH, Shar GQ (2006b). The Status of trace and toxic elements in biol samples (scalp hair) of skin-disease patients and normal subjects. *Turkish J. Med Sci.* 36(4):223-230.
- Ayodele JT, Abubakar MB (1998) Trace element contamination of rain water in the semi arid region of Kano. *Environ. Manag and Health.* 9:176-181.
- Ayodele JT, Abubakar MB (2001). Cleopatra bulimoides and mutellarubens as bio-indicators of trace metals in Tiga Dam, Kano, Nig Res J. *Sci.* 7:45-49
- Ayodele JT, Bayero AS (2002). Arsenic in the blood and urine of some inhabitants of Gombe State, Nig. *ASSET Series B* 1(2):107-117.
- Barrett S (1985). Commercial hair analysis: Sci or scam? *JAMA* 254:1041-1045
- Barnes B, Bradley SG (1994). Planning for a Healthy Baby. Essential Reading for all future parents. Vermillion, London, UK. pp.1-224
- Boeckx RL (1986). Lead Poisoning in Children. *Anal Chem.* 58:274A-288A
- Borel JS, Anderson RA (1984). Chromium in Biochem of the Essent. ultra trace element (Ed: E. Frieden) Plenum Pub. Co. pp. 332 – 45.
- Boris AR (1994). Lead in Hair and Urine of children and Adults from industrialized Areas. *Arch Environ. Health,* 49(1):59 – 62.
- Bord JS, Anderson RA (1984). Chromium in Biochem of the Essent. Ultra trace Elements (Ed: E. Frieden) Plenum Pub. Co
- Bradley SG, Bennett N (1995). Preparation for pregnancy: An Essent. Guide. Argyll Pub. UK. pp. 23-59
- Calabrese EJ (1980). *Nutr. and Environ. Health: The influence of Nutr. Status on Pollutant Toxicity.* Vol1 John Wiley and sons Inc. New York. 600pp.
- Casey CE, Hambidge KM (1980). Trace Element Deficiencies in Man. *Advances in Nutr. Res.* Draper, H.H., ed. Plenum Pub. 3: 55
- Gorecka H, Chojnacki A, Gorecki H (2005). Inter-element interactions in human hair. *Environ. Toxicol. Pharm.* 20:168
- Chojnacki K, Gorecka H, Gorecki H (2006). The influence of living habits and family relationships on element conc. in human hair. *Sci. Total Environ.* 366:612-620.
- DeAntonio SM, Katz SA, Scheimer DM, Wood JD (1982). Anatomically – related variations in trace – metal conc. in hair. *Cli. Chem. Am Assoc. clin. Chem.. Acta* 71: 501 – 10.
- EPA (1979). *Toxic Trace Metals in Mammalian and Human Hair and Nails,* EPA-600/4.049, United States Environ Protection Agency, Research and Development.
- EPA (1980). *Environ. Prot. Agency, Biol. Monitoring of Trace Metals,* EPA600/3-80-089.
- EPA (1988). *Environ. Prot. Agency Special report on ingested inorganic arsenic, skin cancer nutr. Essent.* EPA/625/387/013. July 1988
- Fergusson JE (1990). *The heavy metals: Chem. Environ. Impact and Health Effects.* Pergamon Press, Oxford. 478pp.
- Hambidge KM (1982). Hair analyses: Worthless for vit. limited for minerals .*Am. J. Clin. Nutr.* 36:943-949.
- Hambidge KM Casey CW, Krebs NF (1987). Zinc .In W.Mertz (ed) *Trace elements in human and anim. I nutr.* 5th Edn. Academic Press Inc., San Diego. 2: 50-51.
- Hardling-Barlow I. (1983). In *Arsenic ind. Biomed. Environ. Perspectives* ed Lederer, WH and Fensterim EJ, Van Nostrand Reinhold, New York, 203pp.
- Hill C, Matrone G (1970). Chem. parameters in the studying vivo and invitro interactions of transmissions elements. *Fed. Prod.,* 29: 1474
- Hill GM, Miller ER, Whetter PA, Ultrey DE (1987). Conc. of mineral in tissues, of pigs, from dams fed different levels of dietary zinc. *J. Anim. Sci.* 57:130-138.
- Hopps HC (1977). *The Biol. Bases for using Hair and Nail for Analysis of Trace Elements.* *Sci. Total Environ.* 7: 71-89
- Hove E, Elvehjem CA, Hart EB (1940). The relation of zinc to carbonic anhydrase. *J. Biol. Chem.* 136:425-434.
- Imran H A, Gul KT, Jamali M K, Hassan K, Shar QG (2003). The status of Trace and toxic elements in Biol. samples (Scalp Hair) of skin Disease patients and normal subjects. *Anal chem..*
- Inyengar GV (1984). Preservation and Preparation of Biol. Mater. for Trace Element Analysis: Quality Assurance Considerations, Quality Assurance in Biomedical Neutron Activation Analysis, IAEA Tecdoc-323, Int. Atomic Energy Agency, Vienna 89pp.
- Katz SA (1979). The use of hair as a biopsy mater. for trace elements in the body. *Am. Lab.* 11: 44-52.
- Kazi TG, Jalbani N, Kazi N, Jamali MK, Arain MB, Afridi HI, Kandhro A, Pirzado Z (2008). Eval. of toxic metals in blood and urine samples of chronic renal failure patients, before and after dialysis. *Renal Failure.* 30(7): 737-745.
- Klevay LM (1997). Hair analysis in cli. and experimental med. *Am. J. Cli. Nutr.* 46:233-236
- Kucera J, Lener J, Soukal L, Horakova J (1996). Air Pollution and Biol. Monitoring of Environ. Exposure to Vanadium Using Short Time Neutron Activation Analysis. *Trace Microprobe Techni.* 141(1):193
- Martin RR, Kempson IM, Naftel SJ, Skinner WM (2005). Preliminary synchrotron analysis of lead in hair from a lead smelter worker *Chemosphere* 58: 1385-1390.
- Maugh HT (1978). Hair: A Diagnostic Tool to Compliment Serum and Urine. *Science* 202: 1271-1277.
- Mehra R, Juneja M (2005). Fingernails as bio. indices of metal exposure. *J. Biosci.* 30(2): 101-105.
- Miller ER, Stove HD, Ullrey DE, Ku PK, Hill GM (1979). Copper and zinc in swine nutr. pp.1-139. In *National Feed Ingredients Assoc. lit. review on copper and zinc in anim. nutr.* National Ingredient Assoc. West Des Moines, IA
- Neathery MW, Miller WJ, Blackman DM, Gentry RP, Jones JB (1973). Absorption and tissue zinc content in lactating dairy cows as affected by low dietary zinc. *J. Anim. Sci.* 37:848-852.
- Nnorom IC, Igwe JC, Ejimne JC (2005). Multielement analyses of human scalp hair samples from three distant towns in south eastern Nig. *African J. Biotech.* 4:1124-1127.
- Nowak B (1998). Contents and relationship of elements in human hair for a non-indu. population in Poland. *Sci of the Total Environ.* 209:59-68.
- O'Mahon M (1986). *Sensory eval of food, Stat. methods and procedures.* New York: Marcell Dekker, Inc., pp. 50-55.
- Pirsaraei AS (2007). Lead exposure and hair lead level of workers in a lead refinery ind. in Iran. *Indian J. Occup. Environ.* 11: 6 – 8.
- Potter JL Timmons GD, West R, Silvidi A (1974). Argininosisuccinic acidemia-The Hair Abnormality. *Am. J. Dis Child* 1: 127
- Reinhold JG, Kfoury GA, Ghalambor MA, Jean (1966). Zinc and Copper Conc. in Hair of Iranian Villagers. *Am J. Clin. Nutr.* 18:294-300
- Ryabukin YS (1978). Activation analysis of Hair as an indicator of contamination of man by Environ. Trace element pollutants. IJAEA Report. Vienna: Int. Atomic Energy Agency. IAEA/RL/50.
- Sandra C, Alcides SD, Margarida M, Rejane S, Solange MM, Severino F, Patrick K (2002). Parameters of eval. of Zinc nutr. status. Comparison between zinc hair rates and serum alkaline phosphates in Pre-Schoolers of the Municipality of Joao Pessoa, Paraiba. *Rev. Bras. Saude. Mater infant,* 2(3): 275– 282.
- Scott ML (1972). Trace elements in animal nutr. In J, Mortedt et al (ed) *Micronutrients in agriculture.* SSSA, Madsen, WI. pp. 555-591.
- Strain WH, Steadman LT, Lankau CA, Berlinger WP, Pories, W.J (1996).

- (Analysis of zinc levels in hair for the diagnosis of zinc deficiency in man. *J. Lab. Clin. Med.* 68 (1):244
- Strain, WHS, Pories WJ, Flynn A, Hill OA (1972). Trace Element Nutriture and Metabolism through Head Hair Analysis. In Hemphill, D.D (ed). Trace Substances in Environ. Health Univ. of Missouri Press. Columbia, MO P.383.
- Sukumar A, Subramanian, R (2003). Elements in the hair of non-mining workers of a Lignite open mine in Neyveli. *Ind. Health*, 41: 63 – 8.
- Todd WR, Elvehjem CA, Hart EB (1934). Zinc in the nutrition of the Rat. *Am. J. Physiol.* 107:146-155.
- Tuormaa TE (1994). The adverse effects of lead *J. Nutri Med.* 4(4):351-361
- Tuormaa TE (1995). Adverse effects of zinc deficiency: A review from the lit. *J. Orthomol. Med* 10:149-165.
- Underwood EJ (1977). Trace elements in human and animal nutrition 4th ed. Academic Press. New York 62pp.
- Ulvi H, Yigiter R, Yoldas T, Dolu Y, Var A, Mungen B (2002). Manganese, zinc and copper contents in Hair and their serum conc. in patients with epilepsy. *Eastern J. Med.* 7(2): 3 – 5.
- Vivoli GF, Fantuzzi G, Bergomi M, Tonelli E, Gatto MRA, Zanetti F, Del DM (1990). Relationship Between Zinc in serum and hair and some Hormones During Sexual Maturation in Humans. *Sci. Tot. Environ.* 95:29-40
- Wilhelm M, Hafner D, Lombeck I, Ohnesorge FK (1991). Monitoring of Cadmium, Copper, lead and Zinc status in Young Children using toenails. Comparison with scalp hair. *Sci. Total Environ.* 103 (2 – 3):199 – 207.