

Full Length Research Paper

# Phytochemical and antimicrobial studies of *Commiphora africana* (A. Rich) Engl. root extracts

Akor J. S<sup>1</sup> and Anjorin T. S<sup>2\*</sup>

<sup>1</sup>Department of Chemistry, University of Abuja, Nigeria.

<sup>2</sup>Department of Crop Science, University of Abuja, Nigeria.

Accepted 23 March, 2009

The crude ethanolic extract of *Commiphora africana* root was partitioned with *n*-hexane, chloroform; water and 10% aqueous methanol and screened for chemical constituents and antimicrobial activity. The *n*-hexane fraction, which was the most bio-active was chromatographed on a column silica gel to give a number of purified components. The components from the *n*-hexane were found to be active against *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. *S. aureus* (ATCC 13709) was susceptible to fractions C<sub>1</sub> - C<sub>11</sub> giving zonal inhibition of between 1 - 3 mm. Standard antibiotics - amoxicillin, chloramphenicol and tetracycline were compared with the extracts in antimicrobial activity.

**Key words:** Antimicrobial activity, *Commiphora africana*, phytochemical activity, root extracts.

## INTRODUCTION

*Commiphora africana* (A. Rich) Engl. Syn. *Heudelotia africana* (Family Burseraceae) is a shrub or small tree which has a short lateral branches, sharply pointed at the apex, bearing leaves in small clusters below the tip (Arnold and Dewet, 1993). The plant which is well suited to dry areas is often grown as a hedge in Northern Nigeria (Burkill, 1985). Parts of the plant are medicinally taken in several West African countries possibly because of the presence of phytochemicals such as methylisopropenyl furan, sesquiterpenes and commiphoric acid (Abbiw, 1990).

A macerate of crushed leaves in oil is drunk in Cote d'Ivoire and in Burkina Faso as a sedative and soporific (Adebayo et al., 2006). Bark-extracts have been shown to have some insecticidal activity and to be termite repellent (Abbiw, 1990). The gum is widely used to prepare antiseptic washes and baths for skin infections, sores and leprosy. The seed contains tannin, dye stuff, a fixed oil, dihydroflavonol glucoside and Z-guggulsterone (McGuffin et al., 2006). In Nigeria, a seed decoction is held to be a very effective purgative and vermin-fuge. A dose of 6 g of powdered seed in a glass of water is certain to expel a tapeworm.

In a continued search for new antimicrobial agents from Nigerian higher plants for potential use in medicine

and in crop protection, this work provides a report on the constituents of the biologically active *n*-hexane fraction of the ethanolic crude extract of the root of *C. africana* and was subsequently assessed for their anti-microbial efficacy.

## EXPERIMENTAL

### Materials

The root sample of *C. africana* was collected from Giri village in the Federal Capital Territory, Abuja, Nigeria. This was dried and ground into a coarse powder. Extraction solvents used were ethanol (from BDH Chemicals), ethyl acetate (Rectapur) and *n*-hexane (Merek) and silica gel (Kiesel gel S. 0.2 - 0.5 mm) for column chromatography (from Riedel-Dettaen Ag. Seelze Han-nover). R<sub>f</sub> values were obtained on pre-coated Merek grade TLC plates using a Camag Ultra Violet (U.V) lamp 366 - 254 nm and iodine vapour as detectors.

### Extraction

In the Chemistry Laboratory of University of Abuja, Abuja, Nigeria, the powdered root 400 g was Soxhlet-extracted with 98% ethanol 2.5 L for about 12 h after which the extract was filtered and concentrated to dryness using a rotary evaporator to give 29.5 g of extract.

### Fractions of the crude extract

The extract was dissolved in chloroform 200 ml and taken in a separatory funnel 1 L. The chloroform layer was partitioned with

\*Corresponding author. [oyindamola35@yahoo.com](mailto:oyindamola35@yahoo.com). Tel.: +2348062242866.

water 100 ml x 2, upper aqueous layer separated and concentrated to dryness to obtain dark-brown viscous syrup 13.33 g. The chloroform fraction was further fractionated by first evaporating to dryness, re-dissolved in 10% aqueous methanol 200 ml and extracted with *n*-hexane 200 ml. The two fractions were separated and evaporated to dryness to give *n*-hexane fraction 2.95 g and aqueous methanol fraction 5.20 g.

#### Phytochemical screening of crude extract

Ferric chloride test (Trease and Evans, 1989) on the crude extract gave a blue black coloration on TLC plate. The crude extract also gave positive Lieberman-Burchard test (Finar, 1988)

#### Isolation of constituents of *n*-hexane fraction

##### Vacuum liquid chromatography (VLC)

Vacuum liquid chromatography of the *n*-hexane fraction 2.20 g on TLC grade silica gel as the stationary phase and eluting with mixtures of *n*-hexane and ethyl acetate gave a number of impure fractions (a) 0.10 g (b) 0.01 g (c) 0.23 g (d) 0.09 g (e) 0.05 g (f) 0.16 g and (g) 0.69 g. Fractions c and d were combined together because they contained common components.

##### Column chromatography

Combined fractions c and d 0.32 g were chromatographed over a column of silica gel and eluted with a mixture of *n*-hexane ethyl acetate. 100% *n*-hexane afforded component C<sub>1</sub> (0.02 g); Rf 0.81 (*n*-hexane ethyl acetate 9:1); *n*-hexane ethyl acetate (100:1) gave component C<sub>2</sub> (0.091 g); *n*-hexane ethyl acetate (20:1) afforded component C<sub>3</sub> (0.07 g), Rf 0.63 (*n*-hexane ethyl acetate 9:1). Similar process was carried out on impure fractions (e) to afford C<sub>4</sub> with 100% *n*-hexane 0.017 g, Rf 0.81 (*n*-hexane ethyl acetate 9:1).

The C<sub>4</sub> component was similar to component C<sub>1</sub> obtained from the combined impure fractions c and d and were therefore combined to obtain a total weight of 0.037g. *n*-hexane ethyl acetate 50:1 yielded component C<sub>5</sub> (0.02 g). Rf 0.41 (*n*-hexane: ethyl acetate 9:1). Ethyl acetate 5% in *n*-hexane yielded component C<sub>6</sub> (0.06 g), Rf 0.38 (*n*-hexane: ethyl acetate 9:1).

The impure fraction (g) yielded C<sub>7</sub> with 1% ethyl acetate in *n*-hexane (0.049 g) Rf 0.63 (*n*-hexane: ethyl acetate 1:1). Further elution with 6% ethyl acetate yielded C<sub>8</sub> (0.007 g), Rf 0.72 (*n*-hexane: ethyl acetate 1:1) 8% ethyl acetate in *n*-hexane yielded C<sub>9</sub> (0.00241 g) Rf 0.38 (*n*-hexane: ethyl acetate 1:1). Further elution with 9% ethyl acetate in *n*-hexane yielded C<sub>10</sub> (0.0305 g) Rf 0.39 (*n*-hexane: ethyl acetate 1:1). 15 - 20% ethyl acetate in *n*-hexane yielded C<sub>11</sub> (0.042 g) Rf 0.23 (*n*-hexane: ethyl acetate 1:1).

#### Antimicrobial screening of crude extract and fractions

Crude extract, the *n*-hexane, aqueous and 10% aqueous methanol fractions were screened for antimicrobial activity in the Microbiology Laboratory of National Pharmaceutical Institute, Idu - Abuja, Nigeria, using Agar-Diffusion Technique (Murray, et al., 1995). The extract and fractions (0.01 g each) was added to 5 ml of different solvent to give a concentration of 2,000 µg/ml. One ml of diluted extract was mixed with 19 ml of sterile nutrient agar poured into a sterile Petri-dish and allowed to gel.

The procedure was repeated for each of the test organisms. The inoculated plates were dried and sterile cork borer (No. 4) was used to make four holes evenly distributed in the dried inoculated plates. These holes were filled with the diluted crude extract and fractions.

Positive and negative controls were equally set up. The plates were then incubated at 37 °C for 24 h and the sensitivity results obtained (Table 1).

#### Antimicrobial screening of purified components

The purified components C<sub>1</sub> - C<sub>11</sub> were all screened for antimicrobial activity using a modified version of National Committee for Clinical Laboratory Standards (NCCLS) (Trease and Evans, 1989). Mueller-Hinton agar in molten state 19.8 cm<sup>3</sup> was aseptically incubated with 0.2 cm<sup>3</sup> of each of the test organisms *S. aureus* (ATCC 13709), *E. coli* (NITC 10418), *B. subtilis* (Pharmaceutical Microbiology Dept., ABU, Zaria), *P. aeruginosa* (ATCC 27853) and *C. albicans* (Diagnostic Unit NIPRD, Pharm Micro-biology., Department).

The well mixed media were each dispensed into sterile plates on a flat surface and allowed to gel. Each of the purified components 100 µg was incorporated into sterile disc. Two discs of different components were placed evenly on the surface of each plate, at least 24 mm (centre to centre) between them with the aid of a forceps sterilized via a Bunsen burner. Three standard antibiotic discs (tetracycline, chloramphenicol and amoxicillin) were used as controls. Each of these antibiotic discs was inoculated onto fresh agar/organism surfaces similar to the purified components. The discs gently pressed down onto the Mueller-Hinton agar were left to stand for 15 min. Other controls equally set up were extract, medium and organism controls. The plates in duplicates were incubated at 37 °C for 24 h and the sensitivity results obtained are shown in table 2.

## RESULTS AND DISCUSSION

The crude extract at 2000 µg/ml showed activities against *S. aureus*, *E. coli*, and higher degree of inhibition in *C. albicans* but no activity against *B. subtilis* and *P. aeruginosa* (Table 1). Ten percent methanol aqueous extract showed no activity against *B. subtilis* and *P. aeruginosa* but was active against *S. aureus* and *C. albicans*; and higher degree of inhibition in *E. coli*. The *n*-hexane fraction showed activities against *S. aureus*, *E. coli* and *C. albicans*. Water soluble extract showed activity only on *S. aureus* and *C. albicans*.

Table 2 shows the sensitivity results obtained from the purified components. The *S. aureus* was susceptible to fractions C<sub>1</sub> - C<sub>11</sub> giving zonal inhibition of between 1 - 3 mm. C<sub>1</sub> - C<sub>7</sub> and then C<sub>11</sub>, gave zonal inhibition of 1 mm, C<sub>9</sub>, and C<sub>10</sub>, gave zonal inhibition of 2 mm while C<sub>5</sub>, and C<sub>8</sub>, gave the maximum zonal inhibition of 3 mm. *E. coli* was susceptible to C<sub>1</sub> - C<sub>11</sub> giving 1 - 3 mm zonal inhibition. C<sub>2</sub> - C<sub>9</sub> and C<sub>8</sub> - C<sub>10</sub>, gave 1 mm zonal inhibition, C<sub>1</sub> and C<sub>7</sub> gave 2 mm zonal inhibition and C<sub>6</sub> and C<sub>11</sub> gave 3 mm zonal inhibition. The *B. subtilis* was susceptible to C<sub>8</sub> - C<sub>11</sub> with 1 mm zonal inhibition. *C. albicans* was susceptible to C<sub>6</sub>. *P. aeruginosa* was resistant to all C<sub>1</sub> - C<sub>11</sub>. The controls, OVC (Organism Viability Control) and ESC (Extract Sterility Control) was all positive while the MSC (Media Sterility Control) responded normally.

Standard antibiotics disc tested in parallel with this experiment gave the following results. Amoxicillin showed activity against *C. subtilis*. Chloramphenicol displayed all round activity against all the organisms: 10, 20, 18, 12

**Table 1.** Antimicrobial screening of crude extract and fractions of *C. africana*.

Extracts and Fractions	Micro organisms/Activity				
	<i>S. aureus</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>	<i>C. albicans</i>
Ethanollic Crude extract(2000 µgml <sup>-1</sup> )	+	+	-	-	++
10% aqueous methanol	+	++	-	-	+
Water soluble	+	-	-	-	+
<i>n</i> -hexane	+	+	-	-	+

**Key:** + Activity  
- No activity

**Table 2.** Antimicrobial screening of purified *C. africana* root extract components.

Organism	Component											OVC	S/C	MSC	Chloramphenicol (52 µg)	Amoxillin (25 µg)	Tetracycline (30 µg)
	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	C <sub>10</sub>	C <sub>11</sub>						
<i>C. albicans</i>	0	0	0	0	0	2	0	0	1	0	0	+	NA	N	10	8	8
<i>S. aureus</i>	1	1	1	1	3	1	1	3	2	2	1		NA	O	20	0	12
<i>E. coli</i>	2	1	1	2	1	3	2	1	1	1	3	+	NA	R	18	0	10
<i>B. subtilis</i>	0	0	0	0	0	0	0	1	1	1	1	+	NA	M	12	1	7
<i>P. aeruginosa</i>	0	0	0	0	0	0	0	0	0	0	0	+	NA	A	8	0	0
ESC	+	+	+	+	+	+	+	+	+	+	+	+	NA	L			

**Key:** O/NA = No Activity; S/C = Solvent control, that is, *n*-hexane; ESC = Extract Sterility Control; OVC = Organism Sterility Control; MSC = Medium Sterility Control; + = Normal response.

and 8 mm zonal inhibition against *C. albicans*, *S. aureus*, *E. coli*, *B. subtilis* and *P. aeruginosa* respectively. Tetracyclines also showed relatively all round activity against the five organisms tested for that is 8 mm zonal inhibition for *C. albicans*, 12 mm for *S. aureus*, 10 mm for *E. coli*, 7 mm for *B. subtilis* but no activity against *P. aeruginosa*.

From the above results, the zonal diameter obtained for the purified extracts was  $\geq 3$  mm. Earlier work (Okwute et al., 1989), on the biologically active *n*-hexane extract of the root of *C. africana* showed that the extract contained three triterpenes,  $\alpha$ -amyrin,  $\beta$ -sitosterol and hydroxyl carboxylic acid. The species or strains of the organism could affect the susceptibility or resistivity of the purified extract. This was indicated in the differential response of the tested species to the ethanolic extract components. The C<sub>1</sub> - C<sub>11</sub> fractions were mostly active against *S. aureus* and *E. coli* but least active against *P. aeruginosa* in Mueller-Hinton agar.

Paraskeva et al. (2008) reported from their studies of *in vitro* biological activity of selected South African *Commiphora* species that a greater selectivity was exhibited by the extracts against the Gram-positive bacteria 0.01 - 8.00 mg/ml and the yeasts 0.25 - 8.00 mg/ml than against the Gram-negative bacteria 1.00 - 8.00 mg/ml in an antimicrobial (MIC) assay. Rahman et al. (2008) confirmed the antibacterial efficacy of terpenes from the oleo-resin of *Commiphora molmol* (Engl.). It was found out that higher activity of terpenes 1 - 4 was determined against a multidrug resistant strain of *S. aureus* -SA1199B than other four strains.

## Conclusion

It was confirmed from this experiment that *C. africana* root crude extracts, fractions and components have shown promising but differential *in-vitro* antimicrobial activity. It is possible that more potent components especially against *S. aureus* and *C. albicans* might reside in the polar fractions which should be the subject of future investigation.

## REFERENCES

- Abbiw DK (1990). Useful plants of Ghana. Published by Intermediate Technology Publication Ltd. by S R P, Exeter U K. pp. 232 - 241
- Adebayo AH, Aliyu R, Gatsing K (2006). The effects of *Commiphora africana* (Burseraceae) on serum lipid profile in rats. Int. J. Pharmacol. 2(6): 618-622
- Arnold TH, Dewet S (1993) Plant of Southern Africa: Names and distribution. Mem. Bot. Surv. Africa: 62
- Burkill HM (1985). The Useful Plants of West Tropical Africa, Crown Agents, London, 1, 2nd ed.
- Finar IL (1988). Organic Chemistry (The Lieberman-Burchard Reaction 1885, 1890), 2(5): 518
- McGuffin MJT, Kartesz A, Keung A, Toker, AO (2000). Herbs of commerce, ed. 2 (Herbs commerce ed. 2)
- Murray PR, Ellen B, Pealler MA, Tenover FC, Yolken RH. (1995). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: approved standards- Fifth edition. NCCLS document M7-A5
- Okwute SK, Mitscher LA, Rao SG (1989). Triterpenes from Antimicrobial *C. africana* (Burseraceae) root, J. Chem. Soc. Nigeria, 14: 63-65
- Rahman MM, Garvev M, Pidcock, LJ, Gibbons S (2008). Antibacterial terpenes from the oleo-resin of *Commiphora molmol* (Engl.). Phytother Res. 10:1356-60

Paraskeva MP, Van Vuuren SF, Van Zyl RL, Davids H, Viljoen AM (2008). *In vitro* biological activity of selected South African *Commiphora* species. J. Ethnopharmacol. 119(3):673-9.

Trease GE, Evans WC (1989). A Textbook of Pharmacognosy, 13<sup>th</sup> ed. Bailliere Tinnall Ltd. London

Yamini B, Tripathi OP, Malhatra, SN, Tripathi (1984). Thyroid Stimulating Action of Z-guggulsterone Obtained from *C. mukul*, *Planta Medica*, 78-80