

Full Length Research Paper

# Identification of the S-genotypes of several sweet cherry (*Prunus avium* L.) cultivars by AS-PCR and pollination

Hongxia Wang<sup>1,2</sup>, KaiChun Zhang<sup>3</sup>, Huairui Su<sup>2</sup> and Naihaoye<sup>2, 4\*</sup>

<sup>1</sup>Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China.

<sup>2</sup>College of Horticulture Science and Engineering, Shandong Agriculture University, Tai'an, Shandong 271018, China.

<sup>3</sup>Institute of Forestry and Pomology, Beijing Academy of Agricultural and Forestry Sciences, Beijing 10093, China.

<sup>4</sup>Yellow Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Qingdao 266071, China.

Accepted 30 November, 2009

**Sweet cherry cultivars display a self-incompatibility system that restricts self-fertilization and fertilization between cultivars bearing identical S-alleles. PCR-based S-allele typing system for sweet cherry cultivars has been recently developed. It has been reported that all known self-incompatibility (S) alleles of sweet cherry were more than 16 based on the AS-PCR analysis. In this work, two sets of AS-PCR primers that were designed based on conserved domains of cDNA sequences of S-RNases has been used to characterize the S-genotype of 38 sweet cherry cultivars, including 15 cultivars whose S-genotype had not been previously described. Pollination test was also performed to validate the PCR results, and the consistent results were got. A wide variation in the frequency of S-alleles in the Sweet cherry germplasm was observed. S<sub>3</sub> was the most common in the cultivars evaluated, and S<sub>7</sub>, S<sub>10</sub>-S<sub>16</sub>, S<sub>23</sub>-S<sub>25</sub> as rare allele was not found in China geographical areas in our study.**

**Key words:** Self-incompatibility, *Prunus avium*, S-RNase, AS-PCR, pollination test.

## INTRODUCTION

Correct assignment of sweet cherry cultivars to cross-compatibility groups is important for the efficient production of cherry fruit because most of the sweet cherry cultivars are self-incompatible and inter-incompatible, that is to say, there is no or close to 0% fruit set unless cross-compatible pollination is provided. Self-incompatibility (SI) is a widespread mechanism in flowering plants that prevents self-fertilization and promotes out-crossing. Sweet cherry exhibits stilar monofactorial gametophytic locus self-incompatibility (Crane and Lawrence, 1929). Several works clearly suggest that two different genes of the S-locus control the function of pistil and pollen respectively in GSI (McCubbin and Kao, 2000). The pistil S-gene encodes a family of ribonucleases (S-RNase) (McClure et al., 1989) and a

pollen S-gene candidate encodes an F-box protein, which has recently been described in almond (*Prunus dulcis*) (Ushijima et al., 2003), and in two species of cherry (Yamane et al., 2003). Many worldwide studies on the cross-compatibility of cherry cultivars have been carried on (Crane and Brown, 1937; De Vries, 1968; Tehrani and Brown, 1992).

A classic table assigning some 190 cultivars and a few selections to 13 incompatibility groups and a 'universal donors' group O was established (Matthews and Dow, 1969). Most of the studies above have been done by controlled pollination tests and/or pollen tube growth tests that are time-consuming and tend to be affected by environmental factors. With the finding that S alleles in sweet cherry code for stilar ribonucleases (S-RNase), up to 16 different S-alleles and 27 incompatibility groups have been reported so far using different methods (Tobutt et al., 2001; Sonneveld et al., 2003; Wunsch and Hormaza, 2004a; Boskovic and Tobutt 2001). PCR method using consensus primers, based on conserved

\*Corresponding author. E-mail: [yenh@ysfri.ac.cn](mailto:yenh@ysfri.ac.cn). Tel: 0086 0532 85822957.

**Table 1.** Sweet cherry cultivars used as S-allele standards and their S-genotypes

Cultivar	S-genotype	Publisher (source)
Summit	S <sub>1</sub> S <sub>2</sub>	EM BC NY
Van	S <sub>1</sub> S <sub>3</sub>	EM BC NY
Bing	S <sub>3</sub> S <sub>4</sub>	EM BC NY
Hedelfingen	S <sub>3</sub> S <sub>5</sub>	EM BC NY MI
Governor Wood	S <sub>3</sub> S <sub>6</sub>	EM NY
Burlat	S <sub>3</sub> S <sub>9</sub>	EM BC NY MI

BC: British Columbia (Wiersma et al., 2001).

EM: Easting Malling (Boskovic and Tobutt, 1996; Boskovic and Tobutt, 2001; Boskovic et al., 1997; Sonneveld et al., 2001).

MI: Michigan (Hauck et al., 2001).

NY: New York (Choi. et al., 2000).

**Table 2.** Fragment lengths generated by PCR of specific S-alleles

Allele	DNA fragment length in base pairs for ASPCR	
	Pru-C2m/Pru-C5m	PruT2/SI32
S <sub>1</sub>	886	458
S <sub>2</sub>		422
S <sub>3</sub>	897	306
S <sub>4</sub>	1061	524
S <sub>5</sub>		465
S <sub>6</sub>	560	521
S <sub>9</sub>	780	431

regions (Tao et al., 1999b), or allele-specific primers (Sonneveld et al., 2001, 2003), polymerase chain reaction-restriction fragment length polymorphisms (PCR-RFLPs) (Yamane et al., 2000) or RFLPs (Hauck et al., 2001) has been used in sweet cherry, which is convenient to have a molecular assay that could be used on vegetative material, independently of age or season.

Most sweet cherry cultivars planted by traditional experience in China have not been genotyped yet. In this study, we utilized the PCR-based S-allele typing system to determine the S-genotypes of 38 sweet cherry cultivars, including 15 sweet cherry cultivars whose S-genotype had not been previously described.

## MATERIALS AND METHODS

### Plant material

Young leaf tissue or buds of total 38 sweet cherry cultivars were collected from Institute of Forestry and Pomology, Beijing Academy of Agriculture and Forestry Sciences germplasm garden and Shandong Agriculture University. Six self-incompatibility sweet cherry cultivars with known S-genotype, 'Summit' (S<sub>1</sub>S<sub>2</sub>), 'Van' (S<sub>1</sub>S<sub>3</sub>), 'Bing' (S<sub>3</sub>S<sub>4</sub>), 'Hedelfingen' (S<sub>3</sub>S<sub>5</sub>), 'Governor Wood' (S<sub>3</sub>S<sub>6</sub>) and 'Burlat' (S<sub>3</sub>S<sub>9</sub>) (Table 1) were collected and used as standard

fragment sizes of their corresponding S-alleles in PCR analysis. These cultivars belong to different self-incompatibility groups and their S-genotype have been confirmed in various different works (Wiersma et al., 2001; Boskovic and Tobutt, 1996; Boskovic and Tobutt, 2001; Boskovic et al., 1997; Sonneveld et al., 2001; Hauck et al., 2001; Choi. et al., 2000).

### Isolation of genomic DNA and PCR amplification of S-alleles

Total genomic DNA was isolated from fresh young leaves or buds using the CTAB method described by Stockinger et al. (1996) with a minor modification. The extraction buffer contained 100 mM Tris-HCl (pH 8.0), 20 mM EDTA, 1.4 mM NaCl, 2% CTAB, 2% Polyvinylpyrrolidone (PVP K25) and 1.5% β-mercaptoethanol. The PCR to identify the S-alleles of the different cultivars analysis was carried out using the primer pairs Pru-C2 and Pru-C5 (Tao et al., 1999a) with minor modification, and the combination Pru-T2 (Tao et al., 1999a) and SI32 (Wiersma et al., 2001). Primer sequences were as follows: Pru-C2m (TGGCCAAGTAATTATTCAAACC), Pru-C5m (CAAATACCACTTCATGTAACAAC), Pru-T2 (GTTCTTGCTTTTGCTTTCTTC) and SI32 (CATAGGCCATGGATGGTG). The PCR amplification was carried out in a total volume of 15 μl using 1 μl DNA about 20-30 ng, reagents included 1.5 U of *Taq* DNA polymerase (Promaga), 1×reaction buffer, 0.2 mM dNTP mix, 1.67 mM MgCl<sub>2</sub>, 0.17 μM primer. The samples were run in Thermo cycler (PTC-100; MJ Research). Temperature profiles consisted of an initial denaturing of 3 min at 94 °C, then the samples were cycled 35 times through the following steps: denaturing for 1 min at 94 °C, annealing for 1 min at 56 °C, elongation for 1 min at 72 °C, with a final extension of 10 min at 72 °C. The PCR product S<sub>1</sub>, S<sub>2</sub> of 'Summit'; S<sub>3</sub>, S<sub>4</sub> of 'Bing'; S<sub>5</sub> of 'Hedelfingen'; S<sub>6</sub> of 'Governor Wood'; S<sub>9</sub> of 'Burlat'; S<sub>3</sub> of 'Van' were cloned and sequenced.

### Detection of PCR products


PCR products were run on 4% denaturing PAGE (polyacrylamide gels) containing 4 M urea. Electrophoresis was performed using 1×TBE buffer on a standard thermoplate sequencer gel-electrophoresis unit (BioRad sequi-GenGT Sequencing Cell). Gels were pre-run for 20 min, and then 4 μl samples were loaded into each well. The samples were denatured at 94 °C for 5 min and cool on ice, before loading. Gels were run at 70 W for 3 h, and silver stained for detection.

### Pollination test

Branches bearing buds were cut, stood in water and 'forced' into flower in the lab (about 25 °C). Pollen was collected, dry at 37 °C oven overnight, then 2 mm mesh was used to separate anthers. Flower on branches in the field selected for crossing, emasculation and hand-pollination were done at the balloon stage and then bagged. Each treatment was comprised 50-100 flowers. Compatibility was classified as positive when final fruit set is 5% or more (Way, 1968). Cultivars that were used in pollination test in this research are list in Table 5.

## RESULTS

PCR analysis was conducted using the S-allele consensus primer sets. The known S-genotypes of cultivars represent the S alleles S<sub>1</sub> to S<sub>6</sub> and S<sub>9</sub>. Each of



**Figure 1.** AS-PCR analysis the genotype of different sweet cherry cultivars amplified with primer pair Pru-C2m/Pru-C5m 4,5,7,11,17,24 are the cultivars used as S-genotype standard. \_ emphasize the new S-genotype cultivars.

1 Fengjin (S4S9) 2 Elton heart (S4S6) 3 Hongdeng (S3S9) 4 Governor wood (S3S6) 5 Burlat (S3S9) 6 Moreau (S3S9) 7 Bing (S3S4) 8 Napolen (S3S4) 9 Rainer (S1S4) 10 Celeste (S1S4) 11 Van (S1S3) 12 Hongmi (S3S6) 13 Youyi (S4S9) 14 Yuzhou(S3S9) 15 Stella (S3S4) 16 Zaohongbaoshi (S3S9) 17 Summit(S1S2) 18 Early Rivers (S1S2) 19 Victor (S2S3) 20 Coleney (S5S6) 21 Gil peck (S1S3) 22 Qihao (S1S9) 23 Lg-1 (S1S6) 24 Hedelfigen (S3S5) 25 Jueze (S3S9) 26 Hongmi (S3S6) 27 Juhong (S4S9) 28 Jiahong (S4S6) 29 Lyons (S6S9)

the two pairs of consensus primers amplified one or two bands of various sizes. With the Pru-C2m and Pru-C5m primer set, S<sub>1</sub>, S<sub>3</sub>, S<sub>4</sub>, S<sub>6</sub> and S<sub>9</sub> had the following sizes: 860, 889, 1117, 668 and 806 bp, respectively (Table 2). An example of the amplification obtained with primers Pru-C2m and Pru-C5m is shown in Figure 1. The PruT2 and SI32 primer set was useful for the identification of alleles S<sub>2</sub> and S<sub>5</sub>, which did not amplified with Pru-C2m/Pru-C5m. It was observed that PruT2/SI32 is more efficient since it could amplify all the S alleles we detected. But due to the small size of the fragments that range from approximately 400-550 base pairs, it was not easy to obtain good resolution of the nearly same size bands using 2% standard agarose gels (data not shown). So PAGE (polyacrylamide gels) was used to improve the resolution. Figure 2 shows the high resolution of S<sub>1</sub>-S<sub>6</sub> and S<sub>9</sub> allele with primer set PruT2/SI32.


The S-genotype can be recognized by the comparison of amplified results between the known and unknown genotypes of cultivars. The amplified results in Table 2 were used as the standards to deduce the genotype of others. A total of 14 pollen incompatibility groups in sweet cherry were identified in the 38 cultivars by AS-PCR typing analysis. Among the 38 cultivars used in this work, 20 cultivars were assigned to respective S-allele groups I, II, III, IV, VI, VII, IX and XVI, based on analysis which also matched the previous assignments (Table 3). The genotypes of 18 cultivars that had not been described previously for S-alleles were identified in this work (Table 3). China is not the center of origin of sweet cherry. Therefore, it may be impossible to find the putative new S-alleles in sweet cherry due to germplasm diversity.

However, a new S-genotype combination S<sub>1</sub>S<sub>6</sub> was identified, which should be compatible with cultivars in other groups. Controlled pollination cross was used to validate the genotypes of the new group. The crossing groups of Lg-1 and Juhong are presented in Table 5. The fruit set ratio was used as the criterion of compatibility or incompatibility, and the final fruit set no less than 5% are thought to be compatible. From the results presented in Table 5, we can see that PCR based S-genotype analysis agreed with our controlled pollination data.

The S-genotypes frequency of the 37 cultivars in the 14 groups were different, in which S<sub>3</sub>S<sub>9</sub> had the highest ratio of 22%, that is to say, there are 8 cultivars of S<sub>3</sub>S<sub>9</sub> S-genotypes in the total 37 cultivars S<sub>3</sub>, S<sub>9</sub> are the most-frequent allele; S<sub>1</sub> and S<sub>4</sub> rank the second position (Table 4).

## DISCUSSION

Tao et al. (1999b) developed the AS-PCR typing system for sweet cherry based on the cDNA sequences of S-RNase. This AS-PCR technique has been widely used in apple (*Malus domestica*) (Jassens et al., 1995), Japanese apricot (*Prunus mune*) (Burgos et al., 1998) and almond (*Prunus dulcis*) (Tamura et al., 2000). Designing allele-specific primers are the most important part in this AS-PCR typing system. It has been known that there are five conserved region and two hypervariable regions in the sweet cherry cDNA sequence of the S-RNase (Ishimizu et al., 1998; Tao et al., 1999b). Two sets of primers were designed to anneal the conserved coding region during



**Figure 2.** AS-PCR analysis the genotype of different sweet cherry cultivars amplified with primer pair PruT2/SI32. 2,8,9,11,16,21 are the cultivars used as S-genotype standard. \_ emphasize the new S-genotype cultivars.

1 Coleney (S5S6) 2 Governor wood (S3S6) 3 Jiahong (S4S6) 4 Juhong (S4S9) 5 Hongmi (S3S6) 6 Victor (S2S3) 7 Vega (S2S3) 8 Hedelfigen (S3S5) 9 Burlat(S3S9) 10 Moreau (S3S9) 11 Governor Wood (S3S6) 12 Stella (S3S4) 13 Viva (S2S3) 14,15 Lyons (S6S9) 16 Van(S1S3) 17 Lg-1 (S1S6) 18 Qihao (S1S9) 19 Rainer (S1S4) 20 Gil peck (S1S3) 21 Summit (S1S2) 22 Early Rivers (S1S2).

PCR in our study which spanned the sequence of the two hypervariable regions that usually had intron structures. The intron of each allele differed in sequence and size from all other tested alleles (Tamura et al., 2000). The difference makes the distinguishing of the alleles by PCR possible. However, in some genotypes only one allele could be amplified. Several reasons could be given for this lack of amplification. First, the mismatches between conserved primer sequence and S allele sequence site (Wunsch and Hormaza, 2004a). Second, the S-allele has large intron between the primer pairs. The latter could be the case of allele S<sub>2</sub> that can not be amplified with the primer pair Pru-C2m/Pru-C5m. Third, preferential PCR amplification of some regions while allele could also reportedly be resulting in false negatives (Brace et al., 1993). This work clearly indicates that two PCR reactions with primer pairs Pru-C2m/Pru-C5m and Pru-T2/SI32 are sufficient to identify the S-genotype of sweet cherry cultivars, in most cases due to their high allele polymorphism and high conservation.

Recently the S<sub>7</sub>, S<sub>9</sub>, S<sub>10</sub>, S<sub>11</sub>, S<sub>12</sub>, S<sub>13</sub>, S<sub>23</sub>, S<sub>24</sub> and S<sub>25</sub> S-allele have been identified and the sequences of sweet cherry 10 S-alleles (S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>4</sub>, S<sub>6</sub>, S<sub>9</sub>, S<sub>12</sub>, S<sub>23</sub>, S<sub>24</sub>, S<sub>25</sub>) have been published (Tao et al., 1999a, b; Sonneveld et al., 2001; Wunsch and Hormaza, 2004a). With the increase in number of S-alleles, just primer pairs

Pru-C2m/Pru-C5m and Pru-T2/SI32 may not be enough to differentiate all alleles and identification can be better achieved complementing the results with other conserve primer pairs or with other molecular methods of sweet cherry cultivar identification such as PCR-RFLPs (Yamane et al., 2000). In fact the number of S-alleles in the sweet cherry cultivars that are grown today is limited, as most of these varieties are highly genetically related and there are S-alleles like S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>4</sub>, S<sub>6</sub>, S<sub>9</sub> that are very common among the cultivated varieties (Wunsch and Hormaza, 2004b; Boskovic, 2001). S<sub>7</sub> was first found in the mazzard rootstock 'Charger', then Boskovic found S<sub>7</sub> in two sweet cherry cultivars, 'Gryall's Seedling' and 'Guigne d'Annonay' (Boskovic, 2001). S<sub>10</sub>, S<sub>11</sub> only occurred in the timber clone 'Orleans 171', still have not been found in sweet cherries. Lacis (2008) investigated the S-allele in the Latvian and Swedish germplasm, which appeared to have a high frequency of the S<sub>6</sub> allele in both collections, and a relatively high frequency of the S<sub>5</sub> allele in Latvian germplasm. In this work, S<sub>3</sub>S<sub>9</sub> are the most frequent genotypes. S<sub>3</sub>, and S<sub>9</sub> are the most-frequent allele. S<sub>1</sub> and S<sub>4</sub> alleles rank the second position. The high frequency of S<sub>3</sub> maybe due to its linkage with some important commercial properties, and is that the reason why breeders used materials containing S<sub>3</sub> gene as the parents, or the properties of S<sub>3</sub> are easily

**Table 3.** Incompatibility groups identified by PCR analysis in the sweet cherry cultivars

Incompatibility group	S-genotype	Cultivar
I	S <sub>1</sub> S <sub>2</sub>	Summit Early Rivers
II	S <sub>1</sub> S <sub>3</sub>	Van Gil Peck
III	S <sub>3</sub> S <sub>4</sub>	Bing Napoleon
IV	S <sub>2</sub> S <sub>3</sub>	Stella (sc), Sunburst (sc) Victor Vega Viva
V	S <sub>4</sub> S <sub>5</sub>	-
VI		Governor Wood Hongmi
VII	S <sub>3</sub> S <sub>6</sub>	Hedelfingen
VIII	S <sub>3</sub> S <sub>5</sub>	-
IX	S <sub>2</sub> S <sub>5</sub>	Rainier
	S <sub>1</sub> S <sub>4</sub>	Celeste, Lapins (sc) Viscount
X		Lyons
XI	S <sub>6</sub> S <sub>9</sub>	-
XII	S <sub>2</sub> S <sub>7</sub>	-
XIII	S <sub>6</sub> S <sub>13</sub>	-
XIV	S <sub>2</sub> S <sub>4</sub>	-
XV	S <sub>1</sub> S <sub>5</sub>	Coleney, 6-3
XVI	S <sub>5</sub> S <sub>6</sub>	Burlat, Moreau, Yuzhou,
	S <sub>3</sub> S <sub>9</sub>	Zaohongbaoshi, Jueze, Zaojiang, Hongdeng, Hongyan
XVII		Elton heart, Jiahong
XVIII	S <sub>4</sub> S <sub>6</sub>	Qihao, Zaodaguo
XIX	S <sub>1</sub> S <sub>9</sub>	-
XXI	S <sub>3</sub> S <sub>13</sub>	Fengjin, Youyi, Juhong,
	S <sub>4</sub> S <sub>9</sub>	Changbahong
	S <sub>1</sub> S <sub>6</sub>	Lg-1

The cultivars that had not been described previously are in bold. sc, self-compatible cultivar.

**Table 4.** Frequency of sweet cherry S genotype and allelic frequencies

	S <sub>1</sub>	S <sub>2</sub>	S <sub>3</sub>	S <sub>4</sub>	S <sub>5</sub>	S <sub>6</sub>	S <sub>7</sub>	S <sub>9</sub>	No. alleles	Frequency of alleles
S <sub>1</sub>	-	2	2	3	-	1	-	3	12	0.16
	S <sub>2</sub>	-	3	-	-	-	-	-	5	0.07
		S <sub>3</sub>	-	2	1	2	-	8	20	0.27
			S <sub>4</sub>	-	-	1	-	3	12	0.16
				S <sub>5</sub>	-	2	-	-	3	0.04
					S <sub>6</sub>	-	-	1	7	0.09
						S <sub>7</sub>	-	-	0	0.0
							S <sub>9</sub>	-	15	0.20
74										

**Table 5.** Summarization of controlled pollination cross test results for identification of the S-genotype.

Cultivars	Tester	Fruit set (%)	Compatibility
Lg-1 ( S <sub>1</sub> S <sub>6</sub> )	Governor wood ( S <sub>3</sub> S <sub>6</sub> )	23	+
	Summit ( S <sub>1</sub> S <sub>2</sub> )	24.4	+
	Rainer ( S <sub>1</sub> S <sub>4</sub> )	18.6	+
	Juhong ( S <sub>4</sub> S <sub>9</sub> )	20.2	+
	Elton heart (S <sub>4</sub> S <sub>6</sub> )	22	+
	Van ( S <sub>1</sub> S <sub>3</sub> )	19.4	+
	Bing (S <sub>3</sub> S <sub>4</sub> )	24.3	+
	Moreau (S <sub>3</sub> S <sub>9</sub> )	14.2	+
Juhong ( S <sub>4</sub> S <sub>9</sub> )	Changbahong ( S <sub>4</sub> S <sub>9</sub> )	1.0	-
	Governor wood ( S <sub>3</sub> S <sub>6</sub> )	13.5	+
	Summit ( S <sub>1</sub> S <sub>2</sub> )	13.9	+
	Rainer ( S <sub>1</sub> S <sub>4</sub> )	10.9	+
	Elton heart(S <sub>4</sub> S <sub>6</sub> )	15.2	+
	Van ( S <sub>1</sub> S <sub>3</sub> )	23.8	+
	Bing (S <sub>3</sub> S <sub>4</sub> )	19.5	+
	Moreau (S <sub>3</sub> S <sub>9</sub> )	14.7	+

'-' incompatible. '+' compatible.

to be inherited in the breeding. The sweet cherry cultivars used by Boskovic were mostly from England, American, and Canada which seldom contain S<sub>9</sub>, while most of the cultivars from China and Russia used in our research have proved to have abundant S<sub>9</sub>. And the cultivars we collected rarely have S<sub>2</sub>. The frequency of S-alleles in geographical areas might reflect the local origin of the ancient cultivars.

There are self-fertile cultivars which are derived from crosses obtained through X-irradiated pollen (Lewis, 1949; Lewis and Crowe, 1954) such as 'Stella', 'Lapins', 'Sunburst' etc. Although the band of S<sub>4</sub>' appeared the same as S<sub>4</sub>-allele of other self-incompatibility cultivars in the AS-PCR, it should be possible to select offspring of self-fertile because there is a very tight genetic linkage between the pollen and stilar part genes for Self-incompatibility. So the band S<sub>4</sub>' can be used as a linked marker for the important self-fertile trait in the sweet cherry self-compatibility breeding work. For example offspring yielding the S<sub>a</sub>S<sub>4</sub>' or S<sub>b</sub>S<sub>4</sub>' bands from S<sub>a</sub>S<sub>b</sub> × S<sub>c</sub>S<sub>4</sub>' (c≠4) should be self-compatible, and it also can select offspring of homozygosis S<sub>4</sub>'S<sub>4</sub>' from S<sub>a</sub>S<sub>4</sub>' × S<sub>a</sub>S<sub>4</sub>'. This makes the pre-selection possible, so this technology is promising in the self-compatibility breeding program for its economical and time-saving.

## ACKNOWLEDGEMENTS

This work was supported by the National Natural Science

Foundation of China (Grants # 40706050, 40706048 and 30700619), the National Science and Technology Pillar Program (Grants # 2006BAD01A13, 2008BAC49B04), Qingdao Municipal Science and Technology plan project (Grants # 08-1-7-6-hy) and the Hi-Tech Research and Development Program (863) of China (Grants # 2006AA10Z414).

## REFERENCES

- Boskovic R, Russell K, Tobutt KR (1997). Inheritance of stilar ribonucleases in cherry progenies, and reassignment of incompatibility alleles to two incompatibility groups. *Euphytica* 95: 221-228.
- Boskovic R, Tobutt KR (1996). Correlation of stilar ribonuclease zymograms with incompatibility alleles in sweet cherry. *Euphytica* 90: 245-250.
- Boskovic R, Tobutt KR (2001). Genotyping cherry cultivars assigned to incompatibility groups by analyzing stilar ribonucleases. *Theor. Appl. Genet.* 103: 475-485.
- Brace J, Ockendon DJ, King JG (1993). Identification of S-alleles in *Brassica oleracea*. *Euphytica* 80:229-234.
- Burgos L, Petrez-Tornero O, Ballester J, Olmos E (1998). Detection and inheritance of stilar ribonucleases associated with incompatibility alleles in apricot. *Sex Plant Reprod* 11: 153-158.
- Choi C, Livermor KE, Anderson RL (2000). Sweet cherry pollination : Recombination based on compatibility groups and bloom time. *J. Am. Pomol. Soc.* 54: 148-152.
- Crane MB, Brown AG (1937). Incompatibility and sterility in sweet cherry *Prunus avium* L. *J. Pomol. Hort. Sci.* 15: 86-116.
- Crane MB, Lawrence WJC (1929). Genetical and cytological aspects of incompatibility and sterility in cultivated fruits. *J. Pomol. Hort.* 11: 53-55.

- De Vries DP (1968). Compatibility of cherries in the Netherlands. *Euphytica* 17: 207-215.
- Hauck N, Iezzoni AF, Yamane H, Tao R (2001). Revisiting the S-allele nomenclature in sweet cherry (*Prunus avium*) using RFLP probes. *J. Am. Soc. Hort. Sci.* 126: 654-660.
- Janssens GA, Goderis IJ, Broekaert F, Broothaerts W (1995). A molecular method for S-allele identification in apple based on allele-specific PCR. *Theor. Appl. Genet.* 91: 691-698.
- Lewis D, Crowe LK (1954). The induction of self-fertility in tree fruits. *J. Hort. Sci.* 29: 220-225.
- Lewis D (1949). Structure of the incompatibility gene. II. Induced mutation rate. *Heredity* 3: 339-355.
- Matthews P, Dow KP (1969). Incompatibility groups: sweet cherry (*Prunus avium*). In: Knight, R.L (Eds), Abstract bibliography of fruit breeding and genetics to 1965, *Prunus*, Commonwealth Agricultural Bureaux, Farnham Royal pp. 540-544.
- McClure BA, Haring V, Ebert PR, Anderson MA, Simpson RJ, Sakiyama F, Clarke AE (1989). Style self-incompatibility gene products of *Nicotiana glauca* are ribonucleases. *Nature* 342: 955-957.
- McCubbin AG, Kao TH (2000). Molecular recognition and response in pollen and pistil interactions. *Ann. Rev. Cell Dev. Biol.* 16: 333-364.
- Sonneveld T, Robbins TP, Boškovic R, Tobutt KR (2001). Cloning of six cherry self-incompatibility alleles and development of allele-specific PCR detection. *Theor. Appl. Genet.* 102: 1046-1055.
- Sonneveld T, Tobutt KR, Robbins TP (2003). Allele-specific PCR detection of sweet cherry self-incompatibility (S) alleles S1 to S16 using consensus and allele specific primers. *Theor. Appl. Genet.* 107: 1057-1070.
- Stockinger EJ, Mulinix CA, Long CM, Brettin TS, Iezzoni AF (1996). A linkage map of sweet cherry based on RAPD analysis of a microspore-derived callus culture population. *J. Hered.* 87: 214-218.
- Ishimizu T, Shinkawa T, Sakiyama F, Norioka S (1998). Primary structural features of rosaceous S-Rnase associated with gametophytic self-incompatibility. *Plant Mole. Biol.* 37: 931-1998.
- Tamura M, Ushijima K, Sassa H, Hirano H, Tao R, Gradziel TM, Dandekar AM (2000). Identification of self-incompatibility genotypes of almond by allele-specific PCR analysis. *Theor. Appl. Genet.* 101: 344-349.
- Tao R, Yamane H, Sugiura A, Murayama H, Sassa H, Mori H (1999b). Molecular typing of S-alleles through identification, characterization and cDNA cloning for S-Rnase in sweet cherry. *J. Am. Soc. Hort. Sci.* 124: 224-233.
- Tao R, Yamane H, Sugiura A (1999a). Cloning of genomic DNA sequences encoding S1-, S3-, S4-, and S6-Rnase (accession Nos. AB031815, AB031816, AB031817, and AB031818) from sweet cherry (*Prunus avium* L.). *Plant Physiol.* 121: 1057.
- Tehrani G, Brown SK (1992). Pollen-incompatibility and self-fertility in sweet cherry. *Plant Breed Rev.* 9: 367-388.
- Tobutt KR, Sonneveld T, Boškovic R (2001). Cherry (in)compatibility genotypes-harmonization of recent results from UK, Canada, Japan and USA. *Eucarpia Fruit Breeding Sect. Newsletter* 5: 41-46.
- Ushijima K, Sassa H, Dandekar AM, Gradziel TM, Tao R, Hirano H (2003). Structural and transcriptional analysis of the self-incompatibility locus of almond: identification of a pollen-expressed F-box gene with haplotype-specific polymorphism. *Plant Cell.* 15: 771-781.
- Way R (1968). Pollen incompatibility groups of sweet cherry clones. *Proc. Am. Soc. Hort. Sci.* 92: 119-123.
- Wiersma P, Wu Z, Zhou L, Hampson C, Kappel F (2001). Identification of new self-incompatibility alleles in sweet cherry (*Prunus avium* L.) and clarification of incompatibility groups by PCR and sequencing analysis. *Theor. Appl. Genet.* 102: 700-708.
- Wunsch A, Hormaza JI (2004a). Cloning and characterization of genomic DNA sequences of four self-incompatibility alleles in sweet cherry (*Prunus avium* L.) *Theor. Appl. Genet.* 108: 299-305.
- Wunsch A, Hormaza JI (2004b). Molecular evaluation of genetic diversity and S-allele composition of Spanish local sweet cherry (*Prunus avium* L.) cultivars. *Genet. Res. Crop. Evol.* 51: 635-641.
- Yamane H, Tao R, Murayama H, Sugiura A (2000). Determining the S-genotypes of several sweet cherry cultivars based on PCR-RFLP analysis. *J. Hort. Sci. Biot.* 75: 562-567.
- Yamane H, Ikeda K, Ushijima K, Sassa H, Tao R (2003). A pollen-expressed gene for a novel protein with an F-box motif that is very tightly linked to a gene for S-Rnase in two species of cherry, *Prunus cerasus* and *P. avium*. *Plant Cell Physiol.* 44: 764-769.
- Lacis G, Kaufmane E, Rashal I, Trajkovski V, Iezzoni AF (2008). Identification of self-incompatibility (S) alleles in Latvian and Swedish sweet cherry genetic resources collections by PCR based typing. *Euphytica* 160: 155-163.