

Full Length Research Paper

Hypotensive effects of 3, 4-dihydroxybenzaldehyde isolated from the stem bark of *Musanga cecropioides*.

B.A. Ayinde^{1*}, E.K.I. Omogbai² and D.N. Onwukaeme¹

¹Department of Pharmacognosy, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

²Department of Pharmacology and Toxicology, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

Accepted 12 November, 2009

Previous works on the stem bark of *Musanga cecropioides* R. Brown (Moraceae) reported the hypotensive effects of aqueous extract at 10 - 40 mg/kg. This present work was carried out to examine the hypotensive effects of 3, 4-dihydroxybenzaldehyde isolated from the stem bark. Using pentobarbitone sodium anaesthetized normotensive rabbits; the effects of the compound on blood pressure were tested at doses of 2.5 - 10 mg/kg. Also, the probable mechanism of action of the compound was examined using atropine, promethazine and the direct effect of the compound on isolated heart. At 2.5 mg/kg, the compound lowered the arterial blood pressure by 12.61 ± 2.45 mmHg. The blood pressure was further lowered by 17.88 ± 0.73 mmHg at 10 mg/kg. The hypotensive effect of the compound was observed to be unaffected by prior administration of either atropine (0.5 mg/kg) or promethazine (0.25 mg/kg). Similar to the mechanism of action of the crude aqueous extract, 3, 4-dihydroxybenzaldehyde induced negative inotropic and chronotropic effects on the heart as it dose-dependently reduced the force and rate of contractions of the heart from 0.5 - 2 mg/ml. The reductions observed at 1 mg/ml were effectively reversed by 10 μ g/ml adrenaline. However, the effects of 2 mg/ml were not reversed by administration of the adrenaline. The compound seems to be one of the constituents responsible for the hypotensive effect of the aqueous extract of *M. cecropioides* stem bark earlier reported.

Key words: Hypotensive effects, *Musanga cecropioides*, stem bark, 3, 4-dihydroxybenzaldehyde.

INTRODUCTION

High blood pressure is one of the non-communicable diseases affecting many individuals in both developing and developed countries. When not properly controlled, it could lead to life wasting conditions like stroke, and other cardiovascular complications (Lloyd-Jones et al., 2000). There are various synthetic drugs with different mechanisms of actions, which when appropriately taken as prescribed, have remarkable effects in bringing the condition under control. Effective as these drugs may be, some of them have life threatening side effects, and in some cases they are unavailable to many rural dwellers due to precarious drug distribution and manpower required for the administration of such drugs. Researching into medicinal plants with acclaimed effects in reducing blood pressure therefore becomes imperative in order to

improve the quality of life of the patients. Such attempts will also help examine and validate or otherwise the scientific rationale underlying the consumption of such herbs by people.

Musanga cecropioides R. Brown (Moraceae) has been reported to be one of the plants used in reducing blood pressure in some parts of tropical Africa (Irvine, 1961). This claim, as well as its uses in inducing labour and hypoglycaemic effects, has been scientifically reported on the leaves and the stem bark (Kamanyi et al., 1991, 1992, 1996; Ayinde et al., 2003, 2006; Ade-neye et al., 2007). From phytochemistry point of view, various parts of the plant has been reported to yield some triterpenoid acids like kalaic, musangic, cecro-pioic acid and a host of others (Lontsi et al., 1987, 1989, 1990, 1991a,b, 1992, 1998a,b). In addition, the leaf was reported to contain orientin, isoorientin, vitexin, pro-cyanidins, chlorogenic acid, and catechin that were observed to inhibit angiotensin converting enzyme (Lacaille-Dubois et al., 2001)

*Corresponding author. E-mail: baayinde@uniben.edu.

while we also reported the isolation of 3, 4-dihydroxybenzaldehyde from the stem bark (Ayinde et al., 2007). The compound has been reportedly found in *Pinella ternata* (Thunb.) Breit (Araceae) (Nawalade et al., 2003) and Barley tea in which it was demonstrated to be one of the most potent antioxidant components of the plant (Etoh et al., 2004). The probable cardiovascular effect of the compound is yet to be reported in literature. This work was therefore, carried out to examine the probable effect of this natural product on the blood pressure of normotensive rabbits and also the probable mechanisms of action involved.

MATERIALS AND METHODS

Isolation and characterization of 3, 4-dihydroxybenzaldehyde

The stem bark of *M. cecropioides* was collected in Benin City in May 2004 and authenticated by Mr. Kola Odewo at Forest Research Institute of Nigeria (F.R.I.N., Ibadan) where a herbarium specimen No FHI106428 was deposited. After extraction by decoction method using water, the extract was filtered and concentrated using rotary evaporator as reported earlier (Ayinde et al., 2003). Accelerated gradient chromatographic analysis of the ethyl acetate fraction led to the isolation of the compound with R_f values of 0.48 in $\text{CH}_2\text{Cl}_2\text{-CH}_3\text{OH}$ (19:1) and 0.82 in $\text{CHCl}_3\text{-CH}_3\text{OH}$ (9:1) respectively. The melting point was observed to be 152 - 154°C. Using combinations of nuclear magnetic resonance, mass and ultraviolet spectroscopic analyses, and the compound was characterized as 3, 4-dihydroxybenzaldehyde (Ayinde et al., 2007).

Drugs and chemicals

These include promethazine, pentobarbitone sodium (Sigma Chemicals, England), d-glucose, potassium hydrogen phosphate, magnesium sulphate heptahydrate, calcium chloride dihydrate (Merck, England), atropine (Indus Pharma, India), sodium chloride (BDH Chemicals, England), sodium hydrogen carbonate (T.G.I. Ltd, India), potassium chloride (Cambian Chemicals, England).

Animals

Male matured rabbits were purchased and maintained in the animal house of the Department of Pharmacology and Toxicology with standard animal pellets (Livestock feeds, Benin City, Nigeria) and drinking water *ad libitum* until required. Approval for use of the animals was obtained from the Faculty of Pharmacy Ethical Committee on the use of Animals for Experiments.

Effect of 3, 4-dihydroxybenzaldehyde on the blood pressure

Male rabbits weighing 1.8-2.6 kg were used to test the effects of 3, 4-dihydroxybenzaldehyde on blood pressure. Each animal was anaesthetised with pentobarbitone sodium (40 mg/kg i.p.). After cannulating the ear vein for drug administration, the carotid artery was cannulated and connected via a Bentley Physiological pressure transducer to two-channel Ugo Basile recorder (Gemini, 7070) for recording blood pressure and heart rate. The drug was administered intravenously at doses of 2.5, 5, and 10 mg/kg. A waiting period of 10 min was allowed before the administration of the next dose. The mean \pm SEM in the fall in mean arterial blood

pressure due to each of the doses was obtained from three repetitions.

Experiments on the probable mechanisms of action of the 3, 4-dihydroxybenzaldehyde

Atropine (0.5 mg/kg) was administered into the animal through intravenous route. About 5 min after, 2.5 mg/kg of the 3, 4-dihydroxybenzaldehyde was administered. Similar procedure was followed using promethazine (0.25 mg/kg) (Ayinde et al., 2003).

In order to examine the effects of the compound on isolated rabbit heart, adult rabbits were sacrificed, and the hearts were immediately harvested and placed in dish containing Krebs's physiological solution composed of (mM) -: NaCl 119.0, NaHCO_3 25.0, d-glucose 11.0, KH_2PO_4 1.2, KCl 5.0, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 1.2, CaCl $_2$ 2.6. They were mounted on Langendorff's apparatus connected to a transducer and channel recorder. The hearts were perfused with the physiological solution maintained at 37°C and aerated with 95% O_2 and 5% CO_2 . The set-up was allowed to equilibrate for 20 min before the administration of the compound. The effects of the compound on the isolated heart were tested at 0.5, 1.0 and 2.0 mg/ml. The rate of heart contractions was measured by the number of contractions per minute as indicated on the channel recorder, while the heart contraction force was obtained by comparing the effects of the extract in relation to the deflection produced by a tension of 500 mg set on the transducer before administration of the extract. The experiment was carried out in triplicates.

Statistical analysis

All data were expressed as mean \pm SEM (standard error of mean) and n represents the number of animals used. Where applicable, the data were compared using one way analysis of variance (ANOVA), Graph pad Instant ^R version 2.05a software (UK). The level of significance was from $P < 0.05$.

RESULTS AND DISCUSSION

The 3, 4-dihydroxybenzaldehyde obtained from *M. cecropioides* was observed to affect both the diastolic and systolic blood pressures. The compound exhibited relatively dose dependent reductions in the mean arterial blood pressure of the rabbits used. At 2.5 mg/kg, the compound reduced the mean arterial blood pressure by 12.61 ± 2.45 mmHg and this was further reduced by 17.88 ± 0.73 mmHg at 10 mg/kg (Figure 1). Although there was no significant difference in the reductions produced by 2.5 and 5 mg/kg, however the effects of 2.5 and 10 mg/kg were significantly different ($P < 0.05$).

Prior administration of atropine (0.5 mg/kg) or promethazine (0.25 mg/kg) appeared not to have any effect on the blood pressure reduction activity of the compound as the 2.5 mg/kg of the compound showed similar effect observed in the absence of either the atropine or promethazine (Figure 2). Between 0.5 - 2 mg/ml, the compound produced significant dose related negative inotropic and chronotropic effects on the isolated rabbit hearts. At 0.5 mg/ml, both the force and rate of heart contractions were significantly reduced compared with control levels. The inhibitory effects of the compound at 1 mg/ml were higher




Figure 1. Hypotensive effects of 3, 4-dihydroxybenzaldehyde isolated from the stem bark of *M. cecropioides*. The compound dose dependently reduced blood pressure of the rabbits (n = 3).




Figure 2. The effects of prior administration of either 0.5 mg/kg atropine (upper part) or 0.25 mg/kg promethazine (lower part) on the blood lowering effects of the 3, 4-dihydroxybenzaldehyde. Neither of the two drugs attenuated the effects of the compound on the blood pressure.




Figure 3. Effect 3, 4-dihydroxybenzaldehyde isolated from the stem bark of *M. cecropioides* on the force of contractions of isolated rabbit heart. It remarkably reduced the force of contractions of the heart at 1 and 2 mg/ml. The inhibitory effect of 1mg/ml was reversed by Adrenaline 10 μ g/ml (-----) while the effect of 2mg/ml was irreversible (n = 3).

than those of 0.5 mg/ml while 2 mg/ml exhibited the highest reductions in both the force and rate of contractions. At 1 mg/ml, the inhibitory effects of the compound were effectively reversed by adrenaline 10 μ g/ml. The reversible effects of adrenaline were more pronounced on the heart rate and even higher than the control levels. However, at 2 mg/ml, the adrenaline showed no effect (Figures 2 - 4).

The hypotensive effects of the aqueous extract of the stem bark of *M. cecropioides* were earlier reported. The 3, 4-dihydroxybenzaldehyde reported to be isolated from the stem bark has been established in this work to possess transient but significant dose-related hypotensive properties. The compound was observed to be more active than the crude aqueous extract earlier reported (Ayinde et al., 2003) as it showed activity at a dose of 2.5 mg/kg contrary to the aqueous extract which slightly reduced blood pressure at a dose of 10 mg/kg as reported earlier.

Useful drugs for controlling elevated blood pressure act through various mechanisms. There are those that act by inhibiting angiotensin-converting enzymes, while some have direct effect on the blood vessels by causing remarkable dilatation. Also, there are some hypotensive drugs that have direct effect on the myocardium (Sun et al., 2002). As we reported earlier for this plant, the mechanisms of action of this compound were found to be similar to that of the crude aqueous extract. Like the aqueous extract, the compound's effect on the blood pressure is unaffected by the presence of either atropine or promethazine. This suggests that the activity of the extract did not provoke the release of histamines neither

does it stimulate the muscarinic receptors in the vessels endothelium or the heart. Also, the compound exerts more potent negative inotropic and chronotropic effects on the heart than the crude aqueous extract as reported earlier. This suggests the probable application of the extract and the compound in cases of tachycardia or even arrhythmias. It is possible that the compound reduces the influx of calcium into the myocardial muscles which consequently affects the rate and force of the heart contraction. Adrenaline is a known vasoconstrictor that produces positive inotropic effects on the heart by stimulating adrenoceptors to increase peripheral vascular resistance, reversing peripheral vasodilation. Simple and polyphenolic compounds obtained from some medicinal have been reported to exert remarkable hypotensive. While gallotannin obtained from *Paeonia lactifolia* was reported to be responsible for the blood lowering effect of the plant through its endothelium-dependent vasodilatory effect (Goto et al., 1996), the hypotensive effect of *Dioclea grandiflora* was attributed to dioclein (5, 2'5'-trihydroxy-6, 7-dimethoxyflavone) - which reportedly reduced mean arterial blood pressure via vasodilation in rats by blocking K^+ channels and inducing membrane hyperpolarization (Cortes et al., 2001). Also, Kang et al. (2003) observed that butein (3, 4, 2'4'-tetrahydroxy-chalcone) showed hypotensive effects through endothelium-dependent vasodilation as well as by showing dose-dependent inhibition of angiotensin converting enzyme.

The 3, 4-dihydroxybenzaldehyde isolated from the stem bark of *M. cecropioides* has been established in this work to possess hypotensive effect unhindered by the presence of atropine or promethazine but exerts its effect by




Figure 4. Inhibitory effect of 3, 4-dihydroxybenzaldehyde isolated from the stem bark of *M. cecropioides* on the frequency of contraction of isolated rabbit heart. At 1 and 2 mg/ml it remarkably reduced the rate of contraction of the heart. The inhibitory effect of the sample at 1 mg/ml was reversed by 10 µg/ml Adrenaline (-----). At 2 mg/ml the effect became irreversible (n = 3).




Figure 5. Sample tracings of the effects of 3, 4-dihydroxybenzaldehyde on isolated rabbit heart.

inducing negative inotropic and chronotropic activities on the heart.

ACKNOWLEDGEMENT

The Authors acknowledge the financial assistance of grant obtained from the University of Benin Research and Publication Committee.

REFERENCES

- Adeneye AA, Ajagbonna OP, Ayodele OW (2007). Hypoglycemic and antidiabetic activities of the aqueous and ethanolic extracts of *Musanga cecropioides* in normal and Alloxan induced diabetic rats. *Fitoterapia* 78(7-8): 502-505.
- Ayinde BA, Omogbai EKI, Onwukaeme DN (2003). Pharmacognostic characteristics and hypotensive effect of the stem bark of *Musanga cecropioides* R.Br. (Moraceae). *West Afr. J. Pharmacol. Drug Res.* 19: 37-41.
- Ayinde BA, Onwukaeme DN, Nworgu ZAM (2006). Oxytoxic effects of the water extract on *Musanga cecropioides* R. Brown (Moraceae) stem bark. *Afr. J. Biotechnol.* 5: 1350-1354.
- Ayinde BA, Onwukaeme DN, Omogbai EKI (2007): Isolation and characterization of two phenolic compounds from the stem bark of *Musanga cecropioides* R. Brown (Moraceae). *Acta Pol. Pharm.* 64: 183-185.
- Cortes SF, Rezende BA, Corriu C, Medeiros IA, Teixeira MM, Lomes VS (2001). Pharmacological evidence for the activation of potassium channels as the mechanism involved in the hypotensive and vaso-relaxant effects of dioclein in rat small resistance arteries. *Br. J. Pharmacol.* 133: 849-858.
- Etoh H, Murakami K, Yogohi T, Ishikawa H, Fukuyama Y, Tanaka H (2004): Antioxidant compounds in barley Tea. *BioSci. Biotechnol. Biochem.* 68: 2616-2618.
- Irvine FR (1961): *Wood plants of Ghana*. Oxford University Press, London pp.446-447.
- Goto H, Shimada Y, Akechi Y, Hatori M, Terasawa K (1996). Endothelium –dependent vasodilator effect of extract prepared from the roots of *Paeonia lactifolia* on isolated rat aorta. *Planta Med.* 62: 436-439.
- Kamanyi A, Bopelet M, Aloamaka CP, Obeifuna PCM, Ebeigbe AB (1991). Endothelium – dependent rat aortic relaxation to the aqueous leaf extract of *Musanga cecropioides*. *J. Ethnopharmacol.* 34: 283-286.
- Kamanyi A, Bopelet M, Tatchum TR (1992): Contractile effect of some extracts from the leaves of *Musanga cecropioides* (Cecropiaceae) on uterine smooth muscle of the rat. *Phytother. Res.* 6: 165-167.
- Kamanyi A, Bopelete M, Lontsi D, Noamesi BK (1996): Hypotensive effects of some extracts of the leaves of *Musanga cecropioides* R. Brown (Cecropiaceae) Studies in the cat and the rat. *Phytomed.* 2: 209-212.
- Kang DG, Kim YC, Sohn EJ, Lee YM, Lee AS, Yin MH, Lee HS (2003) Hypotensive effect of butein via the inhibition of angiotensin converting enzyme. *Biol. Pharm Bull.* 26(9): 1345-1347.
- Lacaille-Dubois MA, Frank U, Wagner H (2001): Search for potential Angiotensin converting enzymes (ACE) - inhibitors from Plants. *Phytomed.* 8: 47-52.
- Lloyd-Jones DM, Evans JC, Larson MG, O'Donnell CJ, Rocella EJ, Levy D (2000): Differential control of Systolic and diastolic blood pressure, Factors associated with lack of blood pressure control in the community. *Hypertension* 36: 594-599.
- Lontsi D, Sondengam BL, Ayafor JF (1989): Chemical Studies on the Cecropiaceae – A novel A – ring seco triterpene from *Musanga cecropioides*. *J. Nat. Prod.* 52: 52-56.
- Lontsi D, Sondengam BL, Ayafor JF, Connolly JD (1987). Cecropiacic acid, A new pentacyclic A – Ring Seco triterpenoid from *Musanga cecropioides*. *Tetrahedron Lett.* 28: 6683-6686.
- Lontsi D, Sondengam BL, Ayafor JF, Tsoupras MB, Taracchi R (1990): Further Triterpenoids of *Musanga cecropioides* – the Structure of Cecropic acid. *Planta Med.* 56: 287-289.
- Lontsi D, Sondengam BL, Bodo B, Martin MT (1998a): Kalaic acid, a new Ursane – type Saponin from *Musanga cecropioides*. *Planta Med.* 64: 189-191.
- Lontsi D, Sondengam BL, Bodo B, Martin MT (1998b): Cecropioic acid - a pentacyclic triterpene from *Musanga cecropioides*. *Phytochem.* 48: 171-174.
- Lontsi D, Sondengam BL, Martin MT, Bodo B (1991a). Musancropic and Musancropic acid B-A-ring contracted triterpenes from *Musanga cecropioides*. *Phytochem.* 30: 2361-2364.
- Lontsi D, Sondengam BL, Martin MT, Bodo B (1991b). Seco-ring-A-triterpenoids from the rootwood of *Musanga cecropioides*. *Phytochem.* 30: 1621-1624.
- Lontsi D, Sondengam BL, Martin MT, Bodo B (1992): Musangicic acid, a triterpenoid constituent of *Musanga cecropioides*. *Phytochem.* 31: 4285-4288.
- Nawalade SM, Saagre AP, Lee CY, Kao CL, Tsay HS (2003): Studies on tissue culture of Chinese medicinal plant resources in Taiwan and their sustainable utilization. *Bet. Bull. Acad. Sin.* 44: 79-98.
- Sun J, Khtan B, Huang S, Whiteman M, Zhu Y (2002): Effects of natural products on ischemic heart diseases and cardiovascular system. *Acta Pharmacol. Sin* 23: 1142-1152.