

Full Length Research Paper

Sources of microbial contamination in tissue culture laboratories in southwestern Nigeria

Odutayo, O. I.¹, Amusa, N. A.², Okutade, O. O.¹ and Ogunsanwo Y.R.¹

¹Department of Biological Sciences, Olabisi Onabanjo University, Ago- Iwoye, Nigeria.

Accepted 16 February, 2007

Microbial contamination is a constant problem, which often compromise development of all in vitro techniques. This study aimed at investigating the source of microbial contamination in tissue culture laboratories in southwestern Nigeria. Nineteen microbial contaminants (consisting of eleven bacteria and eight fungi) were found associated with the tissue culture plants and the laboratory environments. The bacterial contaminants include *Pseudomonas fluorescens*, *Escherichia coli*, *Proteus* sp, *Micrococcus* spp, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Bacillus cereus*, *Bacillus subtilis*, *Corynebacterium* sp and *Erwinia* sp. While Fungi isolates were *Alternaria tenuis*, *Aspergillus niger*, *Aspergillus fumigatus*, *Cladosporium* sp, *Saccharomyces* sp, *Fusarium oxysporum*, *Rizopus nigricans* and *Fusarium culmorum*. The rate of occurrence of *S. aureus*, *B. cereus*, *B. subtilis* and *E. coli* were found to be higher (ranging from 36-46%) in human skin than in all other sampled materials. The laboratory walls and tables also harbored most of the contaminating microbes. The laboratory indoor air was found associated with all the contaminating microbes.

Key words: Micro-propagation, plant tissue culture, bacterial and fungal isolates contamination.

INTRODUCTION

The practice of plant tissue culture has contributed towards the propagation of large number of plant from small pieces of stock plant in relatively short period of time (Daniel, 1998). Basically the technique consists of taking a piece of a plant (such as a stem tip, node, meristem, embryo, or even a seed) and placing it in a sterile, (usually gel-based) nutrient medium where it multiplies. In most of the cases the original plant is not destroyed in the process a factor of considerable importance to the owner of a rare or unusual plant. The micro propagation has also been used extensively in the improvement of selections of plant with enhanced stress or pest resistance, production of pathogen free plants and somatic hybridizations (Daniel, 1998). The formulation of the growth medium depends upon whether it is intended to produce undifferentiated callus tissue, multiply the number of plantlets, grow roots, or multiply embryos for "artificial seed.

The nutrient media in which the plant tissue is cultivated is a good source of nutrient for microbial growth. These microbes compete adversely with plant tissue culture for nutrient. The presence of these microbes in these plant cultures usually results in increased culture

mortality, the presence of latent infections can also result in variable growth, tissue necrosis, reduced shoot proliferation and reduced rooting (Kane, 2003).

Although, the tissue culture techniques usually involves growing stock plants in ways that will minimize infection, treating the plant material with disinfecting chemicals to kill superficial microbes and the sterilizing the tool used for dissection, the vessels and media in which cultures are grown (George, 1993). However, contamination has been reported as constant problem, which can compromise development of all in vitro techniques (Enjalric et al., 1988).

About thirty-one micro-organisms from ten different plant cultivars growing in micro-propagation have been isolated identified and characterized, with Yeasts, *Corynebacterium* spp. and *Pseudomonas* spp. being predominant (Leggatt et al., 1994). *Bacillus* sp., *Corynebacterium* sp. and an Actinomycete have also been found contaminating the vitro culture of apple rootstocks (Hennerty et al., 1994). Odutayo et al. (2004) had also reportedly associated the following bacteria *Pseudomonas syringae* pv *phaseolicoli*, *Bacillus licheniformis*, *Bacillus subtilis*, *Corynebacterium* sp and *Erwinia* sp with the contamination of *Hibiscus cannabinus* and *Telfaria occidentalis* in Nigeria.

Therefore since rapid production of pathogen eradica-

*Corresponding author. E-mail: naamusa@softhome.net

ted plants is a fundamental goal of the micro-propagation process, the aim of this research was to investigate and identify sources of microbial contamination of plant tissue cultures in tissue culture laboratories in Nigeria.

MATERIALS AND METHODS

The plant explants used are cassava (*Manihot esculenta*), Kenaf, (*H. cannabinus*) cowpea (*Vigna unguiculata*) and Banana (*Musa paradisiaca*), tissue cultured vessels, the wall and the air in the tissue culture rooms, and transfer rooms and the skin swab of the laboratory staff. The laboratory used includes the tissue culture laboratory at the International Institute of Tropical Agriculture (IITA), that of the Cocoa Research Institute of Nigeria (CRIN), Institute of Agricultural Research and Training (IAR&T) and the Plant quarantine Services (PQS) Headquarter located in Ibadan Nigeria. The plant tissue culture medium used was Murashige and Skoog (1962) medium and sterilized by autoclaving at 121°C for 15 minutes.

Acidified Potato Dextrose Agar (APDA) and Nutrient Agar (NA) were exposed to air in the tissue culture laboratories for a period of 30 s and 60s respectively in each of the laboratories after which the plates were immediately covered and sealed with cellophane. Sterile cotton buds were used to swab 3 cm² on tissue culture walls, tables and body skin of the laboratory staff respectively and kept in sterile bottle.

Sterilization and incubation of plant cultures

The explants were excised and surfaced sterilized by immersion into a 0.75% NaOCl solution for 20 min after rinsing with 70% ethanol for 15 s. The explants were rinsed in 4 successive changes of sterile distilled water. The excised explants were then aseptically transferred to the culture medium, labeled and incubated at 23 ± 1°C during the day and 19 ± 1°C at night for 3 weeks.

Isolation of microbial contaminants

From the contaminated plant tissue culture tubes, emerging microbes were isolated by inoculating them on Acidified Potato Dextrose Agar (APDA) and incubated for 6 days at 26°C under 12 h photoperiod in the case of fungi and on Nutrient Agar incubated for 3 days at 30°C under 12 h photo-period. Pure isolates obtained from repeated sub-culturing of the isolates were placed in an agar slant in MacCarthney bottles and stored at 4°C in a refrigerator.

Characterization and identification of isolates

The fungal isolates were identified using cultural characters and morphology and by comparison with standards (Barnett and Hunter, 1972). In case of bacteria, beside the morphological characteristics, a number of biochemical and physiological tests were carried out on the isolates. The biochemical tests includes Gram staining, spore staining, motility test, catalase production, oxidase test, indole production, citrate utilization, urease activity, Hydrogen sulphide production, gelatin hydrolysis, starch hydrolysis and carbohydrate utilization.

RESULTS AND DISCUSSION

Eighteen microbial contaminants (consisting of eleven bacteria and eight fungi) were found associated with the

tissue culture plants and the laboratory environments (Table 1) The bacterial contaminants includes, *Pseudomonas fluorescens*, *Escherichia coli*, *Proteus sp*, *Micrococcus spp*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Bacillus cereus*, *B. subtilis*, *Corynebacterium sp* and *Erwinia sp*. While Fungi isolates were *Alternaria tenuis*, *Aspergillus niger*, *Aspergillus fumigatus*, *Cladosporium sp*, *Saccharomyces sp*, *Fusarium oxysporum*, *Rizopus nigricans* and *Fusarium culmorum*. The rate of occurrence of bacteria isolates was higher than that of fungal isolates in the plant tissue cultures (Figures 1). *P. fluorescens*, *S. aureus*, *Bacillus cereus*, *B. subtilis*, *Corynebacterium sp* and *Erwinia sp* were found to be the most prevalent on all the sampled plant tissue material However, the rate of occurrence of *S. aureus* in the plant tissue materials was less than 10% (Figure 2).

The rate of occurrence of *S. aureus*, *B. cereus*, *Bacillus reus*, *B. subtilis* and *E. coli* were found to be higher (ranging from 36-46%) in human skin than in all other sampled materials. The laboratory walls and tables also harbored most of the contaminating microbes (Figure 3). The laboratory indoor air was found associated with all the contaminating microbes with the exception of *Erwinia sp*. Microbes are living, biological contaminants that can be transmitted by infected people, animals and indoor air, and they can also travel through the air and get inside homes and buildings. Bacteria species like *Staphylococcus* and *Micrococcus* are found on human skin scales (Trudeau, Fernández-Caldas, 1994). *Pseudomonas*, *Flavobacterium* and *Blastobacter* have been reportedly associated with wet surfaces of air-conditioning systems, cooling coils, drain pans and sump pumps (Trudeau and Fernández-Caldas, 1994). *S. aureus* are emitted from the nasopharynx of normally healthy individuals when the person talks, and are commonly found in air, water, the skin (Trudeau and Fernández-Caldas, 1994).

It was discovered that the microbial population is higher in the preparatory room than the incubating rooms. This might be unconnected with the fact that more people frequent the preparatory room. Flaningan and Morey (1996), reported that presence of bacteria in a room indicate the presence of people and their levels may get high when the building is heavily populated.

Fungal contaminants were also found associated with the indoor air, tables/walls, and human skin (Table 2). Typically, fungi make up two-thirds of all of airborne, living organisms. Miller et al. (1988).had earlier reported *Cladosporium*, *Penicillium*, *Aspergillus* and *Alternaria* as the most common indoor fungi. Regularly used furniture has been reported as a major source of fungal spores (Miller et al., 1988). Marked shade around the house has also been reported to increase indoor fungi counts five-fold (Seltzer, 1995). Fungi grow anywhere indoor, where there is moisture and a food source. Many building materials consist of cellulose materials that are particularly suitable for fungi growth when they are wet. other materials that also support fungi growth include dust, paints,

Table 1. Frequency of occurrence of microbial contaminants in tissue culture laboratories in southwestern Nigeria

	IAR&T/OAU		IITA		CRIN		PQS	
	A	B	A	B	A	B	A	B
<i>Pseudomonas fluorescens</i>	15.67ef	9.33ef	9.00de	5.00ef	13.00de	9.00ef	16.67fg	8.00fgh
<i>Corynebacterium sp</i>	18.33fg	9.00e	10.00e	4.00de	18.00g	8.00def	16.00efg	10.00h
<i>Bacillus subtilis</i>	18.00f	12.00hi	11.00	7.00g	17.00g	8.67ef	13.00def	6.00def
<i>Bacillus cereus</i>	23.67gh	14.00hi	10.67e	6.00fg	14.00ef	9.00ef	14.67efg	9.33gh
<i>Erwinia sp</i>	17.67f	10.00efgh	10.33e	5.67efg	16.00fg	10.00f	17.00g	10.00h
<i>Streptococcus pneumonia</i>	21.33g	10.00efgh	9.33de	4.00de	18.00g	9.00ef	16.00fg	9.00gh
<i>Streptococcus faecalis</i>	18.33fg	11.00h	10.00e	6.00fg	17.00g	7.33cde	12.00cde	5.33cde
<i>Escherichia coli</i>	18.67fg	12.00hi	10.67e	6.00fg	11.00cd	9.00ef	15.67efg	9.00h
<i>Proteus vulgaris</i>	11.00cd	5.00bc	9.00de	5.00ef	10.00bc	8.00def	9.00bc	4.33bcd
<i>Micrococcus sp</i>	13.00de	6.67abcd	9.33de	3.00cd	10.00bc	8.67ef	10.00bcd	3.00bc
<i>Staphylococcus aureus</i>	10.00bcd	6.00bcd	6.67cd	3.00cd	10.00bc	6.00abcd	12.00de	6.33def
<i>Klebsiella aerogenes</i>	6.00a	5.00a	5.00bc	2.00bc	6.30a	5.00abcd	10.67bcd	5.00cde
<i>Alternaria tenuis</i>	7.67a	4.00a	2.00a	0.00a	9.33abc	7.00bcd	10.00bcd	8.00fgh
<i>Aspergillus niger</i>	8.00ab	4.00a	5.00bc	2.00bc	7.67ab	4.67ab	7.67b	4.67bcd
<i>Aspergillus fumigatus</i>	10.33bcd	7.00cd	2.00a	0.00a	7.00ab	3.67a	8.00b	5.33cde
<i>Cladosporium sp</i>	6.00a	6.00abcd	3.33c	4.00de	8.00bc	8.00def	8.00b	5.33cde
<i>Fusarium oxysporium</i>	11.00cd	7.00cd	4.67bc	0.67ab	10.00bc	6.00abcd	7.00b	2.67b
<i>Rhizopus nigricans</i>	7.33ab	4.00a	5.33bc	2.67cd	8.00abc	5.67abcd	2.00a	0.33a
<i>Fusarium culmorum</i>	9.67bc	6.00abcd	5.33bc	2.67cd	6.67a	4.67ab	7.00b	5.33cde
<i>Saccharomyces sp</i>	14.33e	10.00efgh	9.33	5.00ef	14.00ef	6.67bcde	13.67defg	8.00fgh

Figures followed by same alphabet along the columns are not significantly different at 0.05 probability level Using Duncan's Multiple Range Test. A= preparatory room, B= incubating room





Figure 1. Rate of occurrence of microbial contaminant in plant tissue culture

Table 2. The incidence of occurrence of microbial contaminants in Tissue culture laboratory

Microbial contaminants	Plant tissue culture	Human Skin	Laboratory wall/table	Laboratory indoor air	Hand Gloves
<i>Pseudomonas flourescens</i>	35.67f	18.33c	1.00a	5.00ef	13.00de
<i>Corynebacterium sp</i>	28.33e	0.00a	4.00ab	4.00de	18.00g
<i>Bacillus subtilis</i>	25.00d	9.00b	4.00ab	7.00g	17.00g
<i>Bacillus cereus</i>	6.97c	36.00d	9.67b	6.00fg	14.00ef
<i>Erwinia sp</i>	17.67f	4.00a	8.33b	5.67efg	16.00fg
<i>Streptococcus pneumonia</i>	0.00a	31.00d	6.33b	4.00de	18.00g
<i>Streptococcus faecalis</i>	0.00a	21.00c	7.00b	6.00fg	17.00g
<i>Escherichia coli</i>	10.67c	38.00de	12.67bc	6.00fg	11.00cd
<i>Proteus vulgaris</i>	11.00c	8.00b	6.00b	5.00ef	10.00bc
<i>Micrococcus sp</i>	4.00a	18.67c	2.33a	3.00cd	10.00bc
<i>Staphylococcus aureus</i>	0.00a	46.00f	16.67cd	3.00cd	10.00bc
<i>Klebsiella aerogenes</i>	6.00b	0.00a	1.00a	2.00bc	6.30a
<i>Alternaria tenius</i>	12.67c	0.00a	2.00a	0.00a	9.33abc
<i>Aspergillus niger</i>	28.00e	8.00b	5.00ab	2.00bc	7.67ab
<i>Aspergillus fumigatus</i>	21.33d	1.00a	0.00a	0.00a	7.00ab
<i>Cladosporium sp</i>	16.00d	0.00a	3.33a	4.00de	8.00bc
<i>Fusarium oxysporium</i>	21.00d	0.00cd	6.67bc	0.67ab	10.00bc
<i>Rhizopus nigricans</i>	27.33e	4.00ab	5.33ab	2.67cd	8.00abc
<i>Fusarium culmorum</i>	39.67f	1.00a	6.33b	2.67cd	6.67a
<i>Saccharomyces sp</i>	24.33e	6.00b	0.33a	5.00ef	14.00ef

Figures followed by same alphabet along the columns are not significantly different at 0.05 probability level Using Duncan's Multiple Range Test.

**Figure 2.** Incidence of fungal contaminants in plant tissue culture

wallpaper, insulation materials, drywall, grease, soap scum, carpet (especially those backed with jute which is a plant fiber), carpet pads, draperies, fabric, and upholstery (Flannigan and Morey, 1996). Fungi generally need a relative humidity of at least 60% to give them enough moisture to survive or significant moisture intrusion, regardless of humidity.

Sources of indoor moisture that often support fungal

growth includes leaky roofs, damp basement or crawl spaces, house plants watering can generate large amounts of moisture, constant plumbing leaks, carpet directly on cement floors, air-conditioners, drain pans/drip pans under cooling coils (as in refrigerators) and steam from cooking (Flannigan and Morey, 1996)

The microbial contaminant found associated with tissue culture plants includes *P. flourescens*, *Corynebacterium*




Figure 3. The occurrence of bacteria isolates in plant tissue culture

sp, *B. subtilis*, *Proteus vulgaricus*, *Erwinia sp*, *S. aureus*, *E. coli F. culmorum*, *A. niger*, *A. tenius*, *R. nigricans* and *Mucor racemosus*. Odutayo et al. (2004) had earlier reportedly isolated the following contaminants from plant tissue cultures in Nigeria *Pseudomonas syringae* pv *phaseolicoli*, *B. licheniformis*, *B. subtilis*, *Corynebacterium sp* and *Erwinia sp*. While fungal contaminants includes *A. tenius*, *A. niger*, *A. fumigatus* and *F. culmorum*. Leggatt et al. (1994) reported the isolation and characterization of thirty-one microorganisms from ten different plant cultivars growing in micro-propagation, with yeasts, *Corynebacterium spp.* and *Pseudomonas spp.* being predominant. Hennerty (1994) reportedly identified *Bacillus sp.*, a *Corynebacterium sp.* and an Actinomycete as contaminants in the M29 root stocks. Fungal contaminants of plant tissue cultures have also been reported (Kane, 2003). Most of these bacteria contaminants have been reported to increase culture mortality; the presence of latent infections can result in variable growth, tissue

necrosis, reduced shoot proliferation and reduced rooting (Kane, 2003).

Although some of these contaminants might be endogenously embedded in the plant tissues (Pierik, 1988), some might also have emanated from contaminated tools, which were not investigated. Boxus and Terzi (1988) reported that the spread of bacterial contamination was caused by insufficient flaming of contaminated tools and by survival of bacteria in 96% ethanol for a few hours. While flaming for 5 s or more (till the inoculating tools become red hot) did eliminate the spread of bacterial contamination at transfers (Boxus and Terzi, 1987). The use of Bacti-Cinerator during 12 s, by inserting inoculating tools in the middle of the heating element, not on the edges has also proved effective (Singha et al., 1987).

Tissue culture vessels are always closed with loose-fitting caps in order to allow gaseous exchange with the external environment. However, Mites and thrips carrying

fungal spores and bacteria in and on their bodies, often gain entry through this loose fittings and travel from one vessel to another thereby contaminating the cultures. Blake (1994) had earlier reported that fungal contamination of cultures is usually the first sign of a mite or thrip infestation. Hence proper sanitation and effective use of appropriate pesticides to control mites and thrips in tissue culture laboratories will be desirable.

Blake (1994) has reported that thorough disinfections and strict hygiene in the laboratory have achieved effective control of microbial contaminants. Movement of people within the preparatory and incubating rooms in tissue culture laboratory should be reduced significantly to avoid the spread of contaminants. Since bacterial concentrations may be high at both low and high levels of relative humidity; therefore, it is advisable to maintain indoor humidity levels between 40 and 60% (Flannigan and Morey, 1996). Leaked pipes and roofs should be repaired within 24 h of detection; the basement floor should be drained, cleansed and disinfected regularly.

REFERENCES

- Barnett HL, Hunter BB (1972). Illustrated Genera of Imperfect Fungi. Minneapolis: Burgess Publishing Company, Minneapolis MN, p.241.
- Blake J (1994). Mites and Thrips as Bacterial and Fungal Vectors between Plant Tissue Cultures In: Bacterial and Bacteria-like Contaminants of Plant Tissue Cultures ISHS *Acta Horticulturae* 225: <http://www.actahort.org/books/225/index.htm>
- Boxus PH, Terzi JM (1987). Big losses due to bacterial contamination can be avoided in mass propagation schemes. *Acta Horticulturae* 212: 91-93.
- Boxus PH, Terzi JM (1988). Control of Accidental Contaminations during Mass Propagation ISHS *Acta Horticulturae* 225: 198-190.
- Daniel R, Linberge (1998). The many dimension of plant tissue culture research Webmaster of Aggie Horticulture Publications, pp. 201-210.
- Enjalric F, Carron MP, Lardet L (1988). Contamination of Primary Cultures in Tropical Areas: The Case of *Hevea Brasiliensis*. In: Bacterial and Bacteria-like Contaminants of Plant Tissue Cultures ISHS *Acta Horticulturae* 225: [Http://www.actahort.org/books/225/index.htm](http://www.actahort.org/books/225/index.htm).
- Flannigan B, Morey PR (1996) Control of moisture problems affecting biological indoor air quality: International Society of Indoor Air Quality and Climate, Ottawa, Canada, ISIAQ Guideline TF1-1996. <http://www.isiaq.org/>.
- George EF (1993) Plant propagation by tissue culture. Exergetics Ltd., Edington, England. p. 574
- Hennerty MJ, Upton ME, Furlong PA, James DJ, Harris DP, Eaton RA (1994) Microbial Contamination of *In Vitro* Cultures of Apple Rootstocks M26 And M9: In: Bacterial and Bacteria-like Contaminants of Plant Tissue Cultures ISHS *Acta Horticulturae* 225: <http://www.actahort.org/book/225/index.htm>.
- Kane, M (2003) Bacterial and Fungal Indexing of Tissue Cultures <http://www.hos.ufl.edu/mooreweb/TissueCulture/class1/Bacterial%20and%20fungal%20indexing%20of%20tissue%20cultures.doc>.
- Leggatt IV, Waites WM, Leifert C, Nicholas J (1994). Characterisation Of Micro-Organisms Isolated From Plants During Micropropagation In: Bacterial and Bacteria-like Contaminants of Plant Tissue Cultures ISHS *Acta Horticulturae* 225: <http://www.actahort.org/books/225/index.htm>.
- Miller JD (1988). Fungi and fungal products in some Canadian homes. *International Biodeterioration* 24: 103-120.
- Murashige T and Skoog F (1962). A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiologia Plantarum* 15: 475-497.
- Odutayo OI, Oso RT, Akinyemi BO Amusa NA (2004) Microbial contaminants of cultured *Hibiscus cannabalis* and *Telfaria occidentalis* cultured tissue Afr. J. Biotechnol. 3: 301-307.
- Pierik RL (1988) In *Vitro* Cultures of Higher Plants as a Tool in the propagation of Horticultural crops .In: Plant propagation by tissue culture ISHA *Acta Horticulturae*, pp. 24-28.
- Singha S, Bissonette GK, Double ML (1987). Methods for sterilizing Instruments contaminated with *Bacillus sp.* *Hortic. Sci.* 22(4): 659.
- Seltzer JM (1995). Biologic contaminants. *Occupational Medicine: State of the Art Rev.* 10(1):1-25.
- Trudeau WL, Fernández-Caldas E (1994) Identifying and measuring indoor biologic agents. *J. Allergy Clin. Immunol.* 2: 393-400.